

1 **Restoring pre-industrial CO₂ levels while achieving Sustainable Development Goals**

2 **M. E. Capron^{1*}†, J. R. Stewart^{1†}, A. d. R. N'Yeurt^{2†}, M. D. Chambers^{3†}, J. K. Kim⁴, C.**
 3 **Yarish^{5†}, A. T. Jones^{6†}, R. B. Blaylock^{7†}, S. C. James^{8†}, R. Fuhrman^{9†}, M. T. Sherman^{1†}, D.**
 4 **Piper¹, G. Harris¹, M. A. Hasan¹**

5 ¹OceanForesters, Oxnard, CA, USA. ²Pacific Centre for Environment and Sustainable
 6 Development, The University of the South Pacific, Suva, Fiji. ³Department of Food and
 7 Agriculture, University of New Hampshire, Durham, NH, USA. ⁴Department of Marine Science,
 8 Incheon National University, Incheon, South Korea. ⁵Department of Ecology and Evolutionary
 9 Biology, University of Connecticut, Stamford, CT, USA. ⁶Intake Works, Sacramento, CA, USA.
 10 ⁷Thad Cochran Marine Aquaculture Center, University of Southern Mississippi, Ocean Springs,
 11 MS, USA. ⁸Department of Geology, Baylor University, Waco, TX, USA. ⁹Stingray Sensing,
 12 Goleta, CA, USA.

13 *Corresponding author: Mark E. Capron (markcapron@oceanforesters.com)

14 †Part of a MARINER (US Department of Energy ARPA-E) team.

15 **Key Points:**

- 16 • Feeding the world with sustainable seafood from artificial reefs that also restore ocean
 17 biodiversity and support marine protected areas
- 18 • Sustainable development for all while reducing atmospheric CO₂ below 300 ppm by the
 19 2100s and keeping global temperature rise below 1.5°C
- 20 • Quadrupling global electricity production, all carbon-neutral or carbon-negative, while
 21 replacing oil with carbon-negative biofuel
 22

23 **Abstract**

24 A framework is presented with examples of technologies capable of achieving carbon neutrality
 25 while sequestering sufficient CO₂ to ensure global temperature rise less than 1.5°C (after a small
 26 overshoot), then continuing to reduce CO₂ levels to 300 ppm within a century. Two paths bracket
 27 the continuum of opportunities including dry, sustainable, terrestrial biomass (such as
 28 *Miscanthus*, paper, and plastic) and wet biomass (such as macroalgae, food, and green waste).
 29 Suggested paths are adaptable, consistent with concepts of integral ecology, and include holistic,
 30 environmentally friendly technologies. Each path addresses food security, marine plastic waste,
 31 social justice, and UN Sustainable Development Goals. Moreover, oceanic biomass-to-biofuel
 32 production with byproduct CO₂ sequestration simultaneously increases ocean health and
 33 biodiversity. Both paths can accomplish net-zero fossil-CO₂ emissions by 2050. Both paths
 34 include: (1) producing a billion tonnes/yr of seafood; (2) collecting six billion dry tonnes of solid
 35 waste (any mix of organic waste, paper, and plastic) to produce twenty million barrels/day of
 36 biocrude; and (3) installing a million megawatts of CO₂-sequestering (Allam Cycle) electric
 37 power plants initially running on fossil fuels. Resulting food production, solid waste-to-energy,
 38 and fossil-fueled Allam Cycle infrastructure will strengthen the economies in developing
 39 countries. Next steps are (4) sequestering four billion tonnes of byproduct CO₂/yr from solid
 40 waste-to-biofuel by hydrothermal liquefaction; (5) increasing macroalgae-for-biofuel production;
 41 (6) replacing fossil fuel with terrestrial biomass for Allam Cycle power plants; (7) recycling
 42 nutrients for sustainability; and (8) eventually sequestering a total of 28 to 38 billion tonnes/yr of
 43 bio-CO₂ for about \$26/tonne, avoided cost.

44 **Plain Language Summary**

45 Pope Francis, “The urgent challenge to protect our common home includes a concern to bring the
 46 whole human family together to seek a sustainable and integral development...” People can do
 47 this with new ways to grow food from healthy oceans, collect and reuse wastes, make fuel and
 48 electricity, store carbon dioxide, and heal climate change. First, grow more food than needed in
 49 less well-off states, selling some food to better-off states. Second, build ways to reuse the
 50 nutrients and energy in sewage and trash to produce more food and energy. Third, using the
 51 income from food sold to better-off states, install new ways to make energy that address climate
 52 change while making all states better-off. The new energy can start using coal, natural gas, waste
 53 (paper, plastic, food waste), and crop residues, then move to sustainable crops such as
 54 switchgrass, seaweed, and seagrass. The CO₂ normally emitted when making energy is captured
 55 and stored. Growing food and biomass for energy, storing CO₂, and improving ocean
 56 biodiversity and health need to displace the current fossil-fuel industry. These new industries can
 57 work in harmony with all life on Earth for many generations while healing climate change (i.e.,
 58 removing CO₂ from the air).

59 **1 Introduction**

60 This paper presents nations and communities more choices for accomplishing United
 61 Nations Sustainable Development Goals (SDGs) (United Nations, 2016b) while reducing CO₂
 62 levels in ways consistent with integral ecology (Pope Francis, 2015; Sorondo & Ramanathan,
 63 2016) and the UN Agenda for Humanity (United Nations, 2016a). The 2015 Paris Agreement
 64 recommends that “rapid reductions” of greenhouse gases be achieved “on the basis of equity, and
 65 in the context of sustainable development and efforts to eradicate poverty” (IPCC, 2014).

- 66 The 2018 IPCC 1.5°C report (IPCC, 2018) findings include:
- 67 • “Climate-related risks to health, livelihoods, food security, water supply, human security,
 - 68 and economic growth are projected to increase with global warming to 1.5°C and
 - 69 increase further with 2°C.” This implies that paths for addressing climate change will be
 - 70 more easily started and sustained if they improve health, livelihoods, food security, water
 - 71 supply, human security, and economic growth.
 - 72 • “All pathways that limit global warming to 1.5°C with limited or no overshoot project
 - 73 the use of carbon dioxide removal (CDR) on the order of 100–1000 GtCO₂ over the 21st
 - 74 century.” This means that it would be wise to achieve net zero emissions with significant
 - 75 CDR that can be sustained over decades to decrease atmospheric CO₂ concentrations.
 - 76 Simply eliminating fossil fuel use will not keep warming below 1.5°C.
 - 77 • “CDR deployment of several hundreds of GtCO₂ [billion tonnes] is subject to multiple
 - 78 feasibility and sustainability constraints (*high confidence*).” This implies that CDR totals
 - 79 of several thousand billion tonnes of CO₂ require multiple technologies to ensure
 - 80 feasibility and nutrient recycling to ensure sustainability.

81 The UN Environment Programme Emissions Gap Report (UNEP, 2019) “warns that

82 unless global greenhouse gas emissions fall by 7.6 per cent each year between 2020 and 2030,

83 the world will miss the opportunity to get on track towards the 1.5°C temperature goal of the

84 Paris Agreement.”

85 Many authors have looked at a variety of CDR methods (also called negative emissions

86 technologies (NETs)). For example, the U.S. National Academies of Sciences, Engineering, and

87 Medicine (2019) plus the massive literature reviews by Minx et al. (2018), Fuss et al. (2018) and

88 Nemet et al. (2018), as well as Tim Flannery’s books (2015, 2017) and the recent Project

89 Drawdown Review (2020).

90 A big challenge is identifying feasible paths with rapid emission reductions, sufficient

91 CO₂ drawdown, and sustainable at scale for centuries, while also supporting UN Sustainable

92 Development Goals. None of those review publications included three emerging technologies:

93 total ecosystem aquaculture (Capron, 2019; Capron & Piper, 2019; Capron et al., 2018, 2019,

94 2020a, 2020b, 2020c; Chambers, 2013; Lucas et al., 2019a, 2019b; Knowler et al., 2020); Allam

95 Cycle electricity production (8 Rivers Capital, 2020; Allam et al. 2017; Fernandes et al. 2019;

96 McMahon, 2019); and hydrothermal liquefaction (Jiang et al. 2019; Jiao et al. 2017; Pichach

97 2019).

98 Zero carbon electricity such as wind and solar stabilize, but do not reduce CO₂ levels.

99 This paper focuses on reducing CO₂ to pre-industrial levels with emerging carbon negative

100 technologies converting biomass to energy.

101 **2 Methods: Bracketing the choice between bioelectricity and biofuel**

102 The approach of this paper is to solve the climate crisis coincident with achieving SDGs.

103 That is, to solve environmental problems with respect for indigenous cultures (Rockström et al.,

104 2017), all persons, and wildlife while acting to alleviate poverty and prevent armed conflicts


105 (Pope Francis, 2015).

106 This paper presents two paths both of which embody the above goals. The two paths,

107 shown in Fig. 1 as P_{fuel} and P_{electric}, represent extremes of either mostly biofuel or mostly


108 electricity. Two paths are provided so that each community and nation can develop their
 109 individual blend of technology and infrastructure to best fit their unique culture, people, natural
 110 resources, and needs.

111 Both paths have reasonable costs, support sustainable development, get to net zero near
 112 2050, and then sequester legacy CO₂ faster than IPCC pathway P4. This added carbon-removal
 113 capability can overcome potential additional CO₂ releases from land and ocean as the
 114 atmospheric concentrations drop (Keller et al. 2018).



115
 116 Fig. 1. The two paths, P_{fuel} and P_{electric}, superimposed on the four pathways of the IPCC
 117 1.5°C target (IPCC, 2018) reflect annual emissions projections (Figs. SPM.3a and SPM.1b) plus
 118 IPCC cumulative projections (SPM.1c). (Note: Both trajectories between 2020 and 2070 are
 119 based on calculations (SS tabs 1, 2, 10) that connect 2020 emissions with projected post-2070 (or
 120 so) emissions using the technologies presented in this paper.

121 The Supplemental Materials spreadsheet (SS) provides a basis for communities to enter
 122 their own values and calculate scales and costs of food production, energy production, and CO₂
 123 removal from the atmosphere for their path. One pre-calculated path favors dry biomass for
 124 electricity production, the other wet biomass for biofuel production. The proposed technologies
 125 are sufficiently developed to allow accurate estimates of the quantities of products and
 126 byproducts, but less accurate estimates of the costs. One proposed technology (Allam Cycle) will
 127 gasify coal or dry biomass to produce electricity while the other, hydrothermal liquefaction
 128 (HTL), will convert wet biomass into biofuel. Total ecosystem aquaculture (TEA) initially
 129 produces high-value seafood and then scales to produce wet biomass (macroalgae and seagrass).
 130 Calculations are based on Allam Cycle electricity production (8 Rivers Capital, 2020) (SS tabs 9,
 131 12), HTL (Pichach, 2019) (SS tabs 7, 8, 11, 12), and TEA (Capron, 2019; Capron & Piper, 2019;
 132 Capron et al., 2018, 2019, 2020a, 2020b, 2020c; Chambers, 2013; Lucas et al., 2019a, 2019b;
 133 Knowler et al., 2020) (SS tabs 4, 5, 6, 18).



134
135
136
137
138
139

Fig. 2 – Initial technologies to achieve Sustainable Development Goals while restoring 300 ppm CO₂. The gray arrows represent liquid CO₂. Nutrient return from electricity production is dashed because high temperature processes such as gasification, incineration, and pyrolysis, can return some nutrients from the ash, but convert most of the organic nitrogen into nitrogen gas and nitrogen oxides during combustion.

140
141

Both paths achieve SDGs in the sequence shown in Fig. 2, then use the same infrastructure to reduce CO₂ levels (see later sections for details):

142
143
144
145

(1) Install total ecosystem aquaculture (TEA) on floating, flexible, fishing reefs with macroalgae forests to produce seafood while returning nutrients for sustainability (Capron, 2019; Capron & Piper, 2019; Capron et al., 2018, 2019, 2020a, 2020b, 2020c; Chambers, 2013; Lucas et al., 2019a, 2019b; Knowler et al., 2020).

146
147
148
149

(2) Install solid waste-collection systems that pay for themselves by producing bioenergy. Simultaneously install electrical power plants that capture and compress CO₂ normally emitted from coal and natural gas. Switch to capture and compress CO₂ from biomass combustion as rapidly as possible, starting with wastes.

150
151
152
153
154
155

(3) Sequester the captured fossil- and bio-CO₂. Increase the amount of biomass (such as macroalgae, *Miscanthus*, and other sustainable biomass crops) to make carbon negative liquid fossil fuels (using the HTL process which captures some CO₂). Gradually increase the ratio of biomass-fueled electricity to fossil-fueled. Up to 7% of the ocean might be needed for macroalgae (seaweed) to make biofuel on the P_{fuel} path. The rest of the ocean can be marine protected area managed in coordination with indigenous people.

156
157
158

Developed countries might skip the first step while importing high-value seafood from developing countries. Ideally, developed countries lead the second and third steps to more quickly optimize the sustainability and economics of all processes.

159 Table 1 (SS tabs 1, 2) outlines the two alternative global energy demands used to
 160 calculate required bio-CO₂ sequestration by 2070 and beyond. The P_{fuel} path proposes a little
 161 more liquid biofuel than the 2018 demand for oil with relatively little bioelectricity. The P_{electric}
 162 path assumes that the demand for bioelectricity is over twice the 2018 demand for electricity
 163 with one quarter of the 2018 demand for liquid fuel. Specifically, P_{electric} involves mostly electric
 164 transportation, commercial, and residential energy use (little natural gas or biofuels).

165 These two bracketing paths are offered so that each nation and community can plan for
 166 paths within the brackets with the kinds of biomass that fit their resources and needs. Globally,
 167 one path and/or technology is no better than any other; however, at the community level, some
 168 paths and technologies are better than others. At both global and community levels, all paths
 169 satisfy global food demand before significant production of biomass for energy.

170 Table 1 shows estimated plug-in values (in red) and computed numbers (in black). The
 171 variable plug-in numbers are illustrative of possibilities interpolated from 2018 global statistics.
 172 SS tabs 1 and 2 include more plug-in numbers, show the formulae, provide references, and offer
 173 opportunities for various “what if” calculations. The two paths in Tables 1 and 2 are designed to
 174 contrast: (1) P_{fuel}, the “low bioelectricity” path where most electricity is produced by
 175 conventional renewables and nearly all biofuel production is consumed by transportation; and
 176 (2) P_{electric}, the high bioelectricity path that maximizes Allam Cycle bioenergy with carbon
 177 capture and storage (BECCS) with most transportation electrified. P_{fuel} requires somewhat more
 178 biomass production than P_{electric} with a slightly slower return to pre-industrial CO₂ levels.

179 Table 1: Two paths of energy demand and supply a few years after net CO₂ emissions
 180 drop to zero (~2070) (SS tab 1).

Metric	Units	P _{fuel} : low bioelectricity, high biofuel	P _{electric} : high bioelectricity, low biofuel
Global population	Billion	10	10
Projected global average electricity generation in 2070 (2018 world average: 3.5 MWh/capita, China: 5.0, US: 13.6, Japan: 8.3 (BP, 2019))	MWh/yr/person	4	7
Global electricity generation (26 billion MWh/yr in 2018 (BP, 2019))	Billion MWh/yr	40	70
Fraction of global electricity production projected to be BECCS with the remainder nuclear or renewable: solar PV, solar thermal, wind, hydro, wave, geothermal, etc.	%	28%	67%
Global non-electric HTL-produced biofuel use (transportation, industry, heating) (global oil demand of 100 million barrels/day (14 million tonnes/day) or 37 billion barrels/yr in 2018 (U.S. Energy Information Administration, 2020))	Billion barrels/yr	40	10

Global biomass production for Allam Cycle electricity BECCS	Billion dry tonnes/yr	4	17
Global biomass production for non-electric biofuel		35	9
Global biomass production for HTL bio-construction materials (asphalt, plastic, carbon fiber, textiles, etc.)		4	4
Total global biomass production (well past net zero, perhaps 2080)		43	30
Mass of bio-CO₂ captured and stored (well past net zero, perhaps 2080)	Billion tonnes of CO₂/yr	28	38
Year when 2 trillion tonnes of CO₂ are removed from atmosphere and ocean and permanently sequestered	Year	2130	2110


181 Table 2 (SS tab 2) outlines possible approaches to achieve net-zero emissions with some
 182 fossil fuel in 2050 by: (1) capturing and storing all CO₂ emitted by fossil-fuel electricity
 183 generation to make such electricity production carbon neutral, (2) capturing and storing some
 184 CO₂ from biofueled electricity production to offset some non-captured fossil fuel use, (3)
 185 capturing and storing most of the byproduct CO₂ produced when biomass is converted to biofuel
 186 to offset other fossil fuel emissions, and (4) carbon-negative biofuels and electricity replacing
 187 fossil-fueled transportation. Negative emissions from the captured and stored bio-CO₂ offset the
 188 use of fossil fuels (mainly natural gas) for heating and industry. After net zero CO₂ emissions,
 189 increasing biomass-fueled energy production with carbon capture removes CO₂ from the
 190 atmosphere at the rates indicated in Table 1.

191 Table 2: Balancing fossil fuel use, biomass-for-energy production, and bio-CO₂
 192 sequestration for net zero emissions about 2050 (SS tab 2)

Metric	units	Low bio-electricity, high biofuel	High bio-electricity, low biofuel
Global fossil oil and natural gas use without sequestering the CO ₂ .	Billion barrels	29	10
Global negative emissions biofuel production for non-electric use (transportation, industry, heating)	/yr (energy equiv.)	11	0
Global carbon neutral electricity (solar, wind, nuclear, fossil fuel with emissions capture, etc.)	Billion MWh/yr	15	56
Global carbon negative electricity (biomass with carbon capture and sequestration)		25	14
Biomass production at net zero (mix of waste, <i>Miscanthus</i> and similar, and macroalgae)	billion dry tonnes/yr	14	3
Resulting approximation of fossil- and bio-CO ₂	billion	28	28

sequestration (at net zero)	tonnes/yr		
Computed net CO₂ emissions	billion tonnes/yr	0.0	0.0

193 Tables 1 and 2 quantify the steps in Fig. 3 to demonstrate how net-zero emissions are
 194 technically feasible by 2050. Every component of Fig. 3 can scale quickly using existing demand
 195 and supply chains.



196 Fig. 3. Process overview for mature (2070 and beyond) production of food, energy, and
 197 CO₂ sequestration. It is a simplified representation of the future global energy system with future
 198 oceanic integrated food and energy systems. It does not show terrestrial food systems. Each
 199 country and community will determine how much of each component is appropriate depending
 200 on local economics. (Note: *Miscanthus* represents all terrestrial biomass including wood waste,
 201 agricultural residues, etc. that might be gasified directly at the Allam Cycle electricity facility or
 202 fed to HTL. Solid waste represents organic sludges, food waste, paper, and plastics that are not
 203 recycled some other way. Micro- and macroalgae represent all watery biomass including
 204 seagrass and freshwater plants. The darker green and thicker arrows are paths to more bio-CO₂
 205 storage (CO₂ removed from the environment, i.e. negative emissions). The lighter green and
 206 thinner arrows lead to carbon-neutral emissions, including bio-CO₂ emissions from combustion
 207 by airplanes, or wind and solar.
 208

209 The costs, values, and relative local scale for each process and arrow in Fig. 3 can be
 210 modified in the spreadsheet for any given time and location. Potential variations and
 211 uncertainties include how fast will oil prices recover after the COVID-19 pandemic and what are
 212 the effects of millions of barrels per day of inexpensive HTL biocrude made from solid waste?
 213 Price unknowns arise in the early learning curve for employing new technologies. Some carbon
 214 neutral fossil-fueled electricity with 97%+ CO₂ sequestration could continue. Economics are
 215 explained in more detail in the Supplemental Document (SD) and Spreadsheet (SS). Numbered
 216 economic and sustainability considerations labeled in Fig. 3 include:

- 217 (1) Increasing seafood production can start now with excess and artificial nutrients
 218 (subsection 3.1, SS tab 18).
- 219 (2) Ocean (aquatic) plants produce wet biomass feedstock for food and energy (subsection
 220 3.1, SS tabs 4 and 6).
- 221 (3) Wet solid waste is the initial feedstock for HTL biofuel. Dry solid waste can be the initial
 222 biomass feedstock for Allam Cycle electricity.
- 223 (4) About 60% of the carbon in biomass or plastic becomes biocrude oil during HTL
 224 (subsection 3.3). Biocrude can be refined at existing refineries. About 40% of the carbon
 225 can be recovered as a mixture of fuel gas and CO₂ for Allam Cycle (or other) electricity
 226 and heat co-generation or the CO₂ can be separated and sequestered.
- 227 (5) Dry terrestrial biomass can be gasified for Allam Cycle electricity production with carbon
 228 capture and sequestration.
- 229 (6) The Allam Cycle (subsection 3.4) produces electricity from gasified coal, gasified
 230 biomass, or natural gas at 40–60% efficiency while also producing pure CO₂ compressed
 231 to 100-bar ready for sequestration.
- 232 (7) Nutrient recycling is essential for sustained production of seafood and energy.
 233 (subsection 3.2).
- 234 (8) There are many ways to permanently sequester CO₂ (subsection 3.5).

235 **3 Calculations, results, and discussion for each component**

236 3.1 Biomass start-up, eventual scale, cost, and infrastructure

237 Tables 1 and 2 indicate the necessary scale of total biomass production. A higher
 238 proportion of the biomass for the “low bioelectricity, high biofuel” path will be “wet” such as
 239 macroalgae, food and green waste. A higher proportion of the biomass for the “high
 240 bioelectricity, low biofuel” path will be “dry” such as *Miscanthus*, paper and plastic.

241 Wet biomass production starts with seafood grown in total-ecosystem aquaculture (TEA).
 242 The macroalgae biomass grown on flexible floating reef ecosystems (Capron, 2019; Capron &
 243 Piper, 2019; Capron et al., 2018, 2019, 2020a, 2020b; Chambers, 2013; Lucas et al., 2019a,
 244 2019b) and other systems (Buschmann et al., 2017; Kim et al., 2017a, 2017b; Knowler et al.,
 245 2020; Park et al., 2018; Radulovich et al. 2015; Shi et al., 2018) provides food and oxygen for
 246 traditional seafoods (i.e. finfish, crabs, oysters, and the like). Gentry et al. (2017), Froehlich et al.
 247 (2019), and Theuerkauf et al. (2019) provide global overviews of potential locations. The SD
 248 explains TEA adaptation research is needed to ensure seafood and biomass productivity with
 249 biodiversity in warming tropical waters. Fish are currently migrating toward Earth’s poles to
 250 escape marine heat waves (Hastings et al., 2020).

251 While harvesting seafood, macroalgal biomass-for-energy production would be
 252 demonstrated and improved. Fish and shellfish production should cost less than \$2/kg on
 253 average. Domestic sales might be \$1 - \$2/kg while exports earn \$4/kg or more at the dock. At
 254 \$2/kg, a billion wet shell-on tonnes of seafood would be worth \$2 trillion/yr. When demand for
 255 biomass-for-biofuel rises, aquaculture ecosystems can be managed to simultaneously produce
 256 both a billion wet tonnes of seafood and 7 billion wet tonnes (0.7 billion dry tonnes) of

257 macroalgae for energy. At \$100/dry tonne (Lucas et al. 2019a), this start-up macroalgae-for-
 258 energy would be worth \$70 billion/yr.

259 The P_{fuel} path presumes increasing ocean net primary productivity by 40% or about 40
 260 billion dry tonnes/yr. The P_{electric} path projects increasing terrestrial net primary productivity by
 261 15% or about 17 billion dry tonnes/yr. Currently, the world's net primary productivity is near
 262 210 billion tonnes/yr of biomass (Field et al., 1998). Total land productivity is about 110 billion
 263 tonnes on an area of 150 million km². Ocean productivity is about 96 billion tonnes on 360
 264 million km². This implies oceans are under-producing relative to land, which could be remedied
 265 by nutrient recycling and structures supporting macroalgae or seagrass in the photic zone. See
 266 SD Section 3.1 for a discussion of how macroalgae-for-fuel can support biodiversity better than
 267 traditional marine protected areas.

268 Primary conclusions from Table 3 include:

- 269 • There is more-than-sufficient potential additional biomass, between 60 and 100 billion
 270 dry tonnes/yr, much more than the 30–40 billion dry tonnes/yr needed in these
 271 projections. Thus, there is no need to use wood from forests, which is often regarded as
 272 unsustainable (Hudiburg, et al., 2013). More discussion in SD.
- 273 • There is more than enough solid waste (Kaza et al. 2018) for 22 million barrels/day (by
 274 2050) of sweet biocrude oil (SS tabs 4, 7, 11, 12).
- 275 • Every kind of biomass or waste (wet or dry) can contribute, which means every country
 276 can participate in some form of biomass production.
- 277 • While there are obvious differences in maximum scale and cost, most every type of
 278 biomass is significant and is, or will become, a viable industry.
- 279 • These numbers are speculative in that the macroalgae projections are based on theoretical
 280 studies, not physical demonstration projects.

281 Table 3: Estimated global biomass production possibilities for some biomass sources (SS tab 3).

Metric	Estimated global scale at indicated cost ¹	Estimated cost delivered to energy process	Estimated energy-return ratio ²
	billions of dry tonnes/yr	\$/dry tonne	$E_{\text{out}}/E_{\text{in}}$
Organic waste including mixed biosolids, paper, plastic, food waste, etc. ³	5 to 7	\$-200 to \$20	4 to 20
Terrestrial agriculture residues and purpose-grown biomass-for-energy (<i>Miscanthus</i> , etc.) ⁴	6 to 20	\$0 to \$400	1 to 50
Macroalgae with total ecosystem aquaculture ⁵ paying for the structure	0.1 to 0.3	\$ 40 to \$70	20 to 50
Microalgae, mixed species, microbes, and plants ⁶	Small, due to high cost	\$400 to \$2,000	0.4 to 1.1
Macroalgae, anchored systems ⁷	10 to 15	\$125 to \$145	8 to 20
Macroalgae, free-floating systems ⁸	40 to 60	\$75 to \$180	4 to 12

<p>¹ Terrestrial material scale and cost is from references in SS tab 4. Macroalgae scale and cost are interpolated from techno-economic analyses anticipating technologies and systems (SS tab 6). The analyses were funded by the U.S. Department of Energy's Advanced Research Projects-Energy MARINER Program (2017b).</p>
<p>² Terrestrial material energy-return ratios are from references in SS tab 4. Macroalgae energy-return ratios were defined in the MARINER Program as the lower heating value of macroalgae for the energy out (E_{out}) and the non-ambient energy required for planting, growing, harvesting, and transporting to the energy processor for energy in (E_{in}) (ignoring the energy embedded in the structure, ships, etc.). The embedded energy was represented in the capital for ambient energy (wind, wave, solar) in the \$/dry tonne.</p>
<p>³ Solid waste pays a disposal fee as if the HTL unit was a landfill. Landfill fees in the U.S. range from \$30–\$100/wet ton (\$120–\$400/dry tonne) (Environmental Research & Education Foundation, 2019). Negative values (because solid waste pays a disposal fee) could produce oil for \$0/barrel. E_{in} is the difference between the energy expended now to collect and transport solid waste to landfills compared to the energy expended to collect, transport, and process it at HTL facilities. (Quantity from Kaza et al., 2018).</p>
<p>⁴ Based on data from Kaza et al., 2018; Turner et al. 2018; REN21, 2019; U.S. Department of Energy, 2018; Eisentraut, 2010; Daly & Halbleib, 2014; Das et al. 2019; Pandur et al., 2015. (SS tab 4). Significant dry biomass could be delivered to the electricity process (Allam Cycle) for \$50/tonne about the same price as US coal at \$2.5/GJ (\$2.6/MMBTU) (SS tab 4).</p>
<p>⁵ The scale of high-protein products paying for the structure (so that the cost of biomass-for-energy can be as low as \$40/DMT) is limited by humanity's demand for high-protein seafood (Capron & Piper, 2019; Capron et al., 2019, 2020a, 2020b)</p>
<p>⁶ U.S. Department of Energy (2018) and Jiang et al. (2019) projected a range of \$400–\$2,000 per dry ash-free tonne of microalgae in their techno-economic uncertainty analysis. Energy return on investment (EROI) from Zaimes & Khanna (2013).</p>
<p>⁷ The area available for anchored macroalgae systems assumes seafloor depths from 0 to 200-m, generally on relatively flat continental shelves (Lucas et al., 2019a). There are moored systems appropriate for deeper seafloors and steep slopes (Sims et al., 2019). Fig. 11 in SD suggests that wet biomass delivered to the biofuel process (HTL) for less than about \$120/dry tonne would produce biocrude oil for less than about \$70/barrel.</p>
<p>⁸ Free-floating deep-ocean systems access large open-ocean areas by floating in currents, eddies, and gyres with minor steering inputs. Individually free-floating plants include <i>Sargassum</i> (<i>S. fluitans</i> and <i>S. natans</i>) (Sherman et al., 2018). Attached growth plants on free-floating structures (Huesemann et al., 2017) include <i>Saccharina japonica</i>, <i>Saccharina latissima</i>, <i>Undaria pinnatifida</i>, <i>Nereocystis luetkeana</i>, <i>Gracilaria tikvahiae</i>, <i>Gracilaria edulis</i>, <i>Gracilariopsis lemaneiformis</i> and <i>Sargassum polycystum</i>. (SS tabs 4 and 6)</p>

282 The bottom line is that there is more biomass potentially available at reasonable prices
 283 than is needed for either the $P_{electric}$ path, which uses 17 billion dry tonnes of dry biomass for
 284 Allam Cycle and 13 billion dry tonnes of wet biomass (food/green waste + macroalgae) for HTL,
 285 or the P_{fuel} path, which uses 4 billion dry tonnes of dry biomass and 39 billion dry tonnes of wet
 286 biomass (see SS Tabs 2, 3, 4).

287 The availability of large quantities of ocean biomass relieves pressure on terrestrial
 288 sources of biomass, which are increasingly limited by demands for food as well as climate
 289 impacts. TEA could grow 1 billion tonnes/yr of seafood on less than 10% of the suitable

290 continental shelf less than 200-m seafloor depth (identified by Gentry et al., 2017). That would
 291 be about 0.3% of the world's oceans (SS tabs 6 and 18). TEA could grow 39 billion dry tonnes
 292 of oceanic biomass-for-energy on 7% of the world's oceans, including some deep ocean (SD 3.1
 293 and SS tab 6). The remaining 93% of ocean area would not be needed for food or bioenergy
 294 production.

295 3.2 Nutrient recycling start-up, eventual scale, cost, and infrastructure

296 The 17 billion dry tonnes/yr of terrestrial biomass for the P_{electric} path (Table 1) requires
 297 about 50 million tonnes/yr of nitrogen (SS tab 12) and proportional amounts of phosphorous,
 298 potassium, iron, boron, copper, manganese, molybdenum, zinc, nickel, and other micronutrients.
 299 Gasification (start of the Allam Cycle process for coal and dry biomass) converts the nitrogen to
 300 N_2 gas. Lost nitrogen might be made up with advances in nitrogen-fixing crops or increased
 301 artificial nitrogen production. Other nutrients can be recovered from the ash.

302 The 39 billion dry tonnes/yr of oceanic biomass for the P_{fuel} path, requires cycling 1.2
 303 billion tonnes/yr of nitrogen (SS tabs 12 and 18) from the ecosystem-to-energy process and back.
 304 Proportional amounts of phosphorous, potassium, iron, boron, copper, manganese, molybdenum,
 305 zinc, nickel, and other micronutrients cycle along with the nitrogen. HTL recovers virtually all N
 306 as ammonia in the “leftover” water. Other nutrients are recovered in the “ash.” Because recycled
 307 nutrients (such as biosolids) contain a complete array of micronutrients, they are also more
 308 beneficial to biomass growth than commercial fertilizer (Wesseler, 2019; Pan et al., 2017).

309 Other reasons for recycling nutrients (computations and references in SS tab 18):

- 310 • Buying ammonia would add \$24/tonne to the cost of oceanic biomass, (i.e. add US\$22/barrel
 311 to the cost of biocrude oil produced by HTL) based on values used by Jiang et al. (2019).
 312 There are additional costs for other nutrients, such as phosphates.
- 313 • 2018 global artificial nitrogen production of 176 million tonnes of N, production of which
 314 emitted 505 million tonnes of CO_2 . Between 75 and 90% of manufactured ammonia is used
 315 for agriculture. Artificial nitrogen fertilizer production already produces ~1% of global CO_2
 316 emissions (Brown, 2016).
- 317 • If nutrients were not recycled, waste-treatment costs for conventional “wastewater” biologic
 318 nutrient removal processes would increase the cost of bio-oil by \$60/barrel.
- 319 • 1.2 billion tonnes of inorganic nitrogen is available in 2–3 million km^3 of deep ocean water.
 320 Removing the inorganic nitrogen (and other nutrients) from a few million km^3 of deep ocean
 321 water each year is not sustainable. Temporarily using a smaller amount of deep ocean water
 322 to start and expand primary production may be acceptable.
- 323 • Upwelling deep ocean water for nutrient supply brings up CO_2 , drops surface water pH
 324 (ocean acidification), and might increase the amount of CO_2 in the air (Chan et al., 2008;
 325 Chen, 2018; Feely et al., 2008; Köhn et al. 2017; Ries, 2010).
- 326 • Several processes (in addition to HTL, such as anaerobic digestion) convert macroalgae to
 327 energy with good efficiency while separating most of the carbon from the nutrients. These
 328 separated nutrients can be returned to the macroalgae ecosystem during harvesting without
 329 significant cost.

330 3.3 HTL start-up, eventual scale, cost, and infrastructure

331 Recent innovations and cost reductions with HTL (Genifuel, 2019; Jiang et al., 2019; Jiao
 332 et al., 2017; Pichach, 2019; ReNew ELP, 2019; Steeper Energy, 2019; Watson et al., 2019) make
 333 it practical to scale up as a solid waste-collection system that pays for itself. HTL converts any
 334 blend of wet plants, paper, wax, and most plastics to bio-oil – expired juice in plastic bottles,
 335 newspaper, expired packages of meat, seaweed, microalgae, switch grass, feces, biohazard
 336 wastes in plastic – all chopped and blended together. The process is similar to the way algae
 337 became oil when buried deep in the Earth. By using a combination of high temperature (350°C,
 338 660°F) and pressure (200 atmospheres, 3,000 psi) the conversion to oil is complete in about
 339 30 min. Because the reaction temperature is less than 400°C, all plant nutrients can be recovered
 340 and used to grow more plants (see SD 3.3).

341 In the CleanCarbon Energy (CCE) HTL process (Pichach, 2019) about 60% of the carbon
 342 in the biomass becomes biocrude. The other 40% becomes byproduct carbon in the forms of
 343 biochar, CH₄, and CO all of which can be converted to energy and CO₂, which can be captured
 344 for sequestration. SS tab 11 quantifies the amounts of sweet biocrude and the byproduct carbon.

345 HTL technology is nearly commercial now based on substantial research and
 346 development in many countries. Recent examples include work at the U.S. Pacific Northwest
 347 National Laboratories with U.S. Department of Energy funding (Jiang et al., 2019). Aarhus
 348 University (Denmark) has investigated using HTL to recover phosphorus and carbon from
 349 manure and sewage sludge with Horizon 2020 funding (Bruun, 2019). Several companies are
 350 preparing ever larger demonstrations of HTL devices including Genifuel in the USA (2019),
 351 Licella (based in Australia) with a plastic feedstock demonstration in the United Kingdom
 352 (ReNew ELP, 2019), Steeper Energy (2019) in Denmark and Canada, and CleanCarbon Energy
 353 in Canada (Pichach, 2019).

354 Developed countries could accelerate deploying commercial HTL with commercial scale
 355 demonstrations (100 to 4,000 wet tonnes/day). Demonstrations are needed because HTL
 356 processes have been developed so far for less-than-commercial scale with single consistent
 357 feedstocks. Solid waste will be a mixed and inconsistent feedstock requiring more sensors to
 358 predict its properties and controls to produce a consistent refinery-ready biocrude product.
 359 Developed country communities could pay for demonstrations using the disposal fees they
 360 collect to safely recycle and dispose of solid waste. After demonstrations clarify costs, HTL
 361 could be deployed in both developed and developing countries to replace landfills. Each
 362 community would determine their optimum balance between the amount of collected (and
 363 uncollected) waste, their disposal fees, and their resulting income from the sale of biocrude oil,
 364 electricity, and other products. See discussion in Table 3, Note 3, with details and graph in SD.

365 3.4 Allam Cycle start-up, eventual scale, cost, and infrastructure

366 The Allam Cycle (8 Rivers Capital, 2020; Allam et al. 2017; Fernandes et al. 2019;
 367 McMahan, 2019) first makes pure oxygen separated from air. The left-over nitrogen and argon
 368 from air separation can be sold. Inside the Allam Cycle combustion chamber, pure O₂, gasified
 369 coal, gasified biomass or plastic, or natural gas, and CO₂ (for cooling the combustion chamber)
 370 mix. After spinning the turbine, all the CO₂ is compressed and cooled. Most is recirculated. A
 371 little, 3 to 5%, depending on the type of fuel, is available as liquid or supercritical sequestration-

372 ready CO₂. Its pressure, 100 to 150-bar (10 to 15 MPa, 1450 to 2,175 psi), will push it through a
 373 pipeline for direct injection into underground or underwater sequestration.

374 Allam Cycle power plants can produce electricity and byproduct liquid CO₂ using any
 375 biofuel or fossil fuel. Initially, we propose they run on fossil fuels (natural gas or gasified coal)
 376 but be converted to biofuels as rapidly as biofuels become available. Because the fossil-fuel
 377 supply chain and much of the electrical distribution system is already in place, fossil-fueled
 378 Allam Cycle carbon-neutral power plants can replace all expansions and replacements for fossil-
 379 fuel electricity production in less than two decades.

380 There are more designs for electricity with carbon capture and storage than just Allam
 381 Cycle. Several, including Allam Cycle, have detailed technical and cost analyses presented at the
 382 website for the U.S. Department of Energy's Coal FIRST program (2020). Allam Cycle is used
 383 throughout this paper because its projected cost of electricity from coal, with sequestration-ready
 384 CO₂ at 100-bar pressure is \$74/MWh, using typical US coal costs. The other six Coal FIRST
 385 program projects captured a lower fraction of produced CO₂ at 1-bar pressure. Adding \$12/MWh
 386 for compression of CO₂ from 1 to 100-bar (SS tabs 14 and 15), their projected costs ranged from
 387 \$118 to \$243/MWh. (Fuel costs and byproduct sales differ, which complicates this comparison.)

388 James et al. (2019) prepared a standard baseline report for several power plant processes
 389 with CCS. The process with the least avoided cost, supercritical pulverized coal (SC-PC),
 390 levelized cost of electricity is \$64/MWh without CCS or \$109/MWh with 90% carbon capture.
 391 These are James' figures without transport and sequestration (T&S) costs with an added
 392 \$3/MWh to compress from James' 15-bar to Allam Cycles' 100-bar. Irlam (2017) reports similar
 393 values to James. Allam Cycle with CCS in a comparable situation has electricity cost at
 394 \$74/MWh (8 Rivers Capital, 2020). See Section 3.6 for a discussion of costs in terms of \$/tonne
 395 of CO₂ sequestered.

396 8 Rivers Capital (2020) explains that early adopters can sell gas products argon (Ar),
 397 nitrogen (N₂), and CO₂ and use the income to decrease the price of electricity to \$55/MWh (\$54
 398 less than SC-PC coal with CCS).

399 NET Power (a subsidiary of 8 Rivers) targets commercial deployment of 300-MW
 400 natural gas Allam Cycle power plants in 2022 (McMahon, 2019). 8 Rivers has proposed a
 401 demonstration of a 300-MW Allam Cycle with coal gasification at a Wyoming coal mine
 402 including selling all the argon and CO₂. The commercial operation date would be 2026 (8 Rivers
 403 Capital, 2020). Allam Cycle power plants are almost zero emissions and have operating
 404 flexibility that reduces the need for battery backup of solar and wind energy (8 Rivers Capital,
 405 2020). They also provide "firm" power which has been calculated by Sepulveda et al. (2018) to
 406 reduce overall electricity costs in decarbonized scenarios. See discussion in SD Section 3.4.

407 3.5 CO₂ sequestration start-up, eventual scale, cost, and infrastructure

408 There are many options for liquid CO₂ sequestration start-up using the current 13 billion
 409 tonnes/yr of fossil-fueled CO₂ emissions from electricity generation. There are many more
 410 carbon and CO₂ storage techniques appropriate for situations other than low-cost liquid CO₂ not
 411 discussed in this paper. The options shown in Table 4 can retain acceptable costs while scaling
 412 for the safe sequestration of trillions of tonnes of liquid CO₂ produced by the HTL and Allam
 413 Cycle power plants. They include geologic carbon sequestration in depleted oil and gas wells and
 414 brine aquifers (Turner et al., 2018; Deng et al., 2017; Alcalde et al., 2018), basalt and other rocks

415 on land and sub-seafloor (National Academies of Sciences, Engineering, Medicine, 2018;
 416 Kelemen et al., 2019; Snæbjörnsdóttir et al., 2020, Moran et al. 2020), and contained CO₂-
 417 hydrate storage on the seafloor (Brewer, 2000; Capron et al., 2013).

418 SD includes more discussion of the concepts and results in Table 4, including how the
 419 different approaches to CO₂ storage complement each other.

Table 4: Liquid or supercritical CO₂ sequestration scale and cost (SS tab 13).

Metric	Global scale of potential storage ¹	Global scale of injection rate	Cost for injecting into sequestration process with permanent monitoring and occasional repairs ²		Leakage rate ³
	billions of tonnes of CO ₂	billions of tonnes of CO ₂ /yr	US\$/tonne ⁴ of CO ₂	US\$/MWh ⁵ with CO ₂ from Allam Cycle	%
Geologic sequestration ⁶ in emptied oil wells, gas wells, and brine aquifers (negative costs are for enhanced oil recovery (EOR))	2,000 to 5,800	large uncertainty	-\$40 to \$56 most w/o EOR below \$8	-\$27 to \$40, most below \$5	< 0.9% of total per 1,000 years
Mineralization sequestration in on land basalt and peridotite rocks ⁷	more than 1,000	more than 10	\$10 to \$30 on land	\$7 to \$20	Negligible
Mineralization sequestration in subsea basalt rocks ⁷	more than 20,000	much more than 20	\$200 to \$400	\$140 to \$300	Negligible
Contained CO ₂ -hydrate storage on the seafloor ⁸	more than 20,000	much more than 20	\$5 to \$10	\$3 to \$7	<0.06% per 1,000 years
¹ Many countries have the resources for only one of the four options. Not every option is sure of the necessary scale.					
² The cost range for geologic storage represents variations in geology, meaning some countries will have inexpensive storage sites and some will have expensive geologic storage. Mineralization costs depend on the characteristics of the local rocks and the depth of drilling required. The range for hydrate storage costs reflects the current situation of relatively little research and development.					
³ Leakage of 0.9% over 1,000 years (Alcade et al., 2018) applied to 2 trillion tonnes of CO ₂ would be 18 million tonnes of CO ₂ /yr. Or 0.06% over 1,000 years (Capron et al., 2013) applied to 2 trillion tonnes is only 1 million tonnes of CO ₂ /yr. (SS tab 13).					
⁴ Costs do not include capturing, compressing, and transporting pure CO ₂ (projected at \$1/t in tab 14 of Supplemental Spreadsheet). Transportation costs are highly dependent on distance to suitable storage location estimated at \$2 to \$3/t for 100 km (National Academies of Sciences, Engineering, Medicine, 2019).					

<p>⁵ Different fuels have different \$/MWh (with the same \$/tonne of CO₂) due to differences in their electrical efficiency and their carbon:hydrogen ratio. This column shows the \$/MWh using gasified coal into Allam Cycle plant. The Supplemental spreadsheet shows it for other fuels. Note that US\$10/MWh corresponds to 1 cent/kWh</p>
<p>⁶ With geologic or mineralization storage, the injection rate of CO₂ should not exceed that which causes earthquakes or leaks due to high pressure in the ground near the injection point (Deng et al., 2017).</p>
<p>⁷ The actual mineralization rate depends on the characteristics of the local rocks (McGrail et al., 2017; Snæbjörnsdóttir et al., 2020). See SD for maps and discussion of different types of rocks with more references (Gunnarsson et al., 2018; Kelemen et al., 2019; Moran et al., 2020).</p>
<p>⁸ Contained ocean hydrate storage scale and injection rate is essentially unlimited. It may be the least expensive option for coastal communities with small continental shelves. From Capron et al. (2013) but updated in SS tab 17.</p>

420 3.6 Paying for removing and storing legacy CO₂

421 Legacy CO₂ is commonly thought of as CO₂ from emissions already in the atmosphere
 422 and ocean (Friedlingstein et al., 2019; Knutti & Rogelj, 2015). Our calculation includes future
 423 fossil-fuel CO₂ uncaptured emissions in the total legacy CO₂ to be removed from the air and
 424 oceans. The total cost is a cost to society in the form of higher energy costs. The cost calculation
 425 below is an apples-to-apples comparison with:

- 426 ○ \$150/tonne for direct air capture (in 2019 USD) (Baker et al., 2020);
- 427 ○ \$74/tonne (\$52/MWh) (James et al., (2019) breakeven emissions penalty (aka
 428 “avoided”) cost when adding CCS to a SC-PC coal power plant, the lowest cost
 429 option in James’ Exhibit ES-4). Costs may be slightly higher when biomass replaces
 430 coal (BECCS).

431 3.6.1 Capture

432 The first added cost and energy component of removing and storing legacy CO₂ is for
 433 capture. That is concentrating the CO₂ about 2,500 times from a little over 0.04% in air to >95%.
 434 Allam Cycle power plants always capture the CO₂ when they produce electricity, so the added
 435 cost for capture is zero.

436 3.6.2 Compression

437 The second added cost and energy component is for compressing the pure CO₂ to a liquid
 438 or supercritical state for permanent sequestration, which varies for the following different
 439 situations:

- 440 • CO₂ capture from Allam Cycle – Each 300-MW coal- (and likely biomass)-fired power
 441 plant compresses 4,600 tonnes/hr of CO₂ from 30 to 150 bar. Most of the CO₂ is
 442 recirculated working fluid. About 230 tonnes/hour is produced from coal for sale or
 443 sequestration. The energy required to compress CO₂ from a gas at 30 bar to a
 444 supercritical fluid at 100 to 150 bar is small, about 9 kWh/tonne (8 Rivers Capital, 2020).
 445 The combined energy plus other operating and capital costs are near \$1/tonne of CO₂ for
 446 coal or \$2/tonne for natural gas. This is based on data from Fernandes et al. (2019), Atlas
 447 Copco CO₂ compressors (2020), Allam et al. (2017), and 8 Rivers Capital (2020) (SS tab
 448 14).

- 449
- 450 • CO₂ capture from HTL – HTL produces bio-crude plus fuel gas that could be combusted
 - 451 with air such that it produces gas with a high fraction of CO₂ (10 to 20%) at 1 bar.
 - 452 Capturing >95% of the CO₂ costs about \$40/tonne of CO₂. Compressing CO₂ from 1 to
 - 453 100 bar requires about 130 kWh/tonne of CO₂. The combined capture, energy plus other
 - 454 operating, and capital costs are near \$65/tonne of CO₂. Most of the cost is for capture and
 - 455 compressing energy, which varies significantly by location, by technology, and over
 - 456 time, as indicated in Table 5.
 - 457 • Hybrid of HTL co-located with Allam Cycle – HTL's byproduct fuel gas and CO₂ at 1
 - 458 bar would be blended and provided as fuel (low-grade fuel gas) to the Allam Cycle. Its
 - 459 value as fuel should cover the cost of compressing it to the required fuel pressure. This
 - 460 situation's capture and compression cost should be similar to the \$1 or \$2/tonne of CO₂

461 3.6.3 Transportation

462 The third added cost component (relatively little energy needed because the CO₂ is a

463 supercritical fluid with very little friction) is the capital cost for transportation, which has been

464 projected by National Academies of Sciences, Engineering, Medicine (2019) as \$2/tonne for a

465 100 km pipeline.

466 3.6.4 Storage

467 The fourth added cost is sequestration of the pure, compressed CO₂. Table 5 (SS tabs 13

468 and 14) values are based on transportation and storage costs of \$10/tonne of CO₂. (We note that

469 James et al. (2019) and Rubin (2015) used \$9/MWh in SC-PC avoided cost calculations (\$12 per

470 tonne of CO₂) for transportation and storage.) This paper used \$8/tonne of CO₂ as an average

471 cost of sequestering liquid or supercritical CO₂ because Turner et al. (2018), Deng et al. (2017)

472 and others project costs for many saline aquifers as \$1 - \$8/tonne. In addition, Table 4 shows

473 negative costs (a credit) for those able to sell CO₂ for EOR (see more discussion in SD).

474 3.6.5 Input fuel cost

475 A fifth cost component is the varying cost of fuels plus economics of the new and old

476 technologies for converting fuel into liquid fuel and/or electricity and process heat. For example,

477 if the new fuel source is less expensive (such as solid waste) than the old fuel (such as liquified

478 natural gas), capturing and sequestering CO₂ might have negative additional cost (Table 5, first

479 two rows). Similarly, if the new fuel source costs \$11/GJ (such as HTL biocrude from

480 macroalgae) instead of \$2.5/GJ (U.S. coal), the additional cost might be \$180/tonne of CO₂

481 (Table 5, bottom row).

482 3.6.6 Process cost

483 A sixth cost component is because different processes result in different leveled
 484 electricity cost (\$/MWh) even with the same fuel cost (\$/MMBTU). The process cost may also
 485 be expressed in \$/tonne of CO₂ captured and compressed. The mainstream processes competing
 486 with Allam Cycle for fossil fuel or biomass electricity are supercritical pulverized coal (SC-PC)
 487 and combined cycle gas turbine (CCGT using natural gas). The Allam Cycle process cost with
 488 CCS appears to be \$15/tonne of CO₂ higher than for SC-PC without CCS (based on statements in
 489 8 Rivers Capital (2020), explained in SD, used in Table 5).

490 3.6.7 Total cost

491 Each row in Table 5 presents the sum of the six cost components to society of producing
 492 electricity, capturing CO₂, compressing it to liquid, transporting it, and permanently sequestering
 493 it, while showing the outcomes using the Allam Cycle process and varying fuel cost.

494 The transportation and sequestration cost of \$10/tonne of CO₂ is included in all rows.
 495 (Rows 1 and 2 are negative because the cost is offset by income from waste disposal fees and
 496 sales of gases.) A local analysis is required to show the local cost differences for each technology
 497 with the local cost of fuel. The assumptions and variables in Table 5 include (see calculations
 498 and explanations in SS tab 14):

- 499 1) Waste can be converted to inexpensive energy with CO₂ capture and sequestration
 500 because disposal fees decrease the cost of fuel.
- 501 2) Terrestrial (dry) biomass (agricultural wastes and purpose-grown biomass) costs about
 502 the same as coal. That might be \$1.9/GJ in some countries (such as US) and \$4.7/GJ in
 503 other countries (such as Japan which is dependent on imported coal at about \$100/tonne).
- 504 3) The hybrid of HTL co-located with Allam Cycle has about the same added cost for
 505 sequestering CO₂ as does Allam Cycle alone (greatly reducing the sequestration cost for
 506 the byproduct fuel gas and CO₂ generated during HTL).
- 507 4) HTL biocrude and biogas made from purpose-grown biomass are likely to cost much
 508 more than coal or natural gas as shown in the bottom two rows of Table 5. Therefore, we
 509 assume essentially no HTL biocrude-from-macroalgae will be fed into Allam Cycle
 510 plants for electricity production; it will be used for transportation fuels.

SS tab 16 includes a traditional calculation of “avoided” or “breakeven emissions penalty” costs. With SC-PC_{ref} and Allam Cycle_{CCS} the avoided cost is \$22/tonne of CO₂. This compares well with the slightly more conservative \$26/tonne of CO₂ shown in Table 5.

Table 5. Added cost to society for capturing, compressing, and sequestering CO₂, changing from various fossil fuels to biomass fuels, and changing to Allam Cycle. Each row reflects a different local situation. Negative numbers mean reduced costs but are limited to early adopters using dry waste for fuel and able to sell gases. “No gas sales” means demand for more CO₂, argon, or nitrogen has dropped to zero.

Metric	Additional \$/tonne of CO ₂	Comment
Allam Cycle power plant gasifying \$0/GJ	-\$250	Lower electricity fuel cost

(\$0/MMBTU) dry waste in place of \$7.6/GJ (\$8/MMBTU) LNG, including income from sales of argon and nitrogen plus CO ₂ for EOR		possible when retaining solid waste disposal fees to offset Allam capital and operating costs.
Allam Cycle power plant gasifying \$0/GJ (\$0/MMBTU) dry waste in place of \$2.5/GJ Illinois coal delivered in US, including income from gas sales	-\$34	
Allam Cycle power plant burning terrestrial biomass delivered for the same \$2.5/GJ as for US coal, no gas sales	\$26	When fuel costs the same, all the additional cost is process change (\$15/tonne), compressing (\$1/tonne), transporting, and sequestering liquid CO ₂ (\$10/tonne).
Allam Cycle power plant burning \$11/GJ HTL biocrude instead of fossil oil for the same \$11/GJ, no gas sales	\$26	
Hybrid co-located HTL and fossil-fired (some HTL biogas) Allam Cycle capturing and compressing CO ₂ from both processes. Same \$/GJ for biomass or fossil fuel, no gas sales	\$26	
Standalone HTL facility using by-product biogas internally with internal capture and compression of by-product CO ₂ , no gas sales	\$75	Using historic capture and compression average cost of \$65/tonne plus the same \$10/tonne for sequestration.
Allam Cycle power plant burning \$11/GJ HTL biocrude in place of \$7.6/GJ LNG (approximate), no gas sales	\$90	Higher fuel cost increasing electricity price is most of the added expense.
Allam Cycle power plant burning \$11/GJ HTL biocrude in place of \$2.5/GJ coal (approximate), no gas sales	\$180	

Table 2 shows about 28 billion tonnes/yr of fossil- and bio-CO₂ being sequestered on either path at net zero emissions. With mostly co-located HTL and Allam Cycle facilities, the global cost is 28 billion tonnes/yr times \$26/tonne, which rounds to \$730 billion/yr.

511 A range of 28 to 38 billion tonnes/yr of bio-CO₂ is being sequestered in Table 1 on either
 512 path for reducing atmospheric CO₂ concentrations (carbon dioxide removal (CDR)). Suppose an
 513 additional 20 billion tonnes/yr of fossil-CO₂ is generated and sequestered. The average net mass
 514 sequestered between the two paths is 53 billion tonnes times \$26/tonne (from Table 5), which
 515 rounds to \$1,400 billion/yr with mostly co-located HTL and Allam Cycle facilities.

516 If HTL is not co-located with Allam Cycle facilities, both paths would use \$75/tonne for
 517 HTL byproduct CO₂ capture, compression, and sequestration. The HTL-focused P_{fuel} path would
 518 cost about \$2,300 billion/yr. The Allam Cycle-focused P_{electric} path would total about \$1,900
 519 billion/yr (SS tab 14).

520 US\$1,400 billion/yr is \$175/person/year for 8 billion people, \$700/yr for a family of four
 521 (much better than CDR at \$150/tonne, which would cost a family of four nearly \$4,000/yr). On
 522 the other hand, \$1,400 billion is only 1.6% of the total global 2019 gross domestic product of
 523 \$87 trillion (StatisticsTimes, 2019). The SD provides more discussion about the following:

- 524 • Process cost explained
- 525 • Putting the cost of sequestering CO₂ in perspective
- 526 • Lower costs for early adopters
- 527 • Allocating costs for removing legacy CO₂
- 528 • Examples of fossil-CO₂ fees and sequestration payments

529 3.7 SDGs path

530 These multiple interrelated systems can start by achieving UN SDGs and expand in scale
 531 to reduce CO₂ levels. These systems are interrelated in that the most circular economy (cradle-to-
 532 cradle manufacturing) and the best economics occur when the systems are co-located. Systems
 533 include the following:

534 3.7.1 Food systems

535 Total ecosystem aquaculture systems are built-reef ecosystems with nutrient recycling
 536 that can provide abundant, inexpensive multi-species seafood. Distributed globally, seafood reefs
 537 (Capron, 2019; Capron & Piper, 2019; Capron et al., 2018, 2019, 2020a, 2020b; Chambers,
 538 2013; Lucas et al., 2019a, 2019b) can sustainably and economically produce a billion tonnes/yr
 539 of seafood, 300 grams/person/day for 8 billion people. (The FAO (2018) estimates current
 540 seafood production (including aquaculture and wild-caught) near 170 million tonnes/yr.)
 541 Developing countries might earn income from developed countries initially by exporting
 542 seafood. Developing countries might earn income from developed countries by accommodating
 543 refugees and migrants as temporary or permanent guest workers on their built-reef ecosystems.
 544 Aquatic-based organic fertilizers can replace chemical fertilizers. Scaling built-reef total
 545 ecosystem aquaculture allows more marine protected areas.

546 3.7.2 Human waste resource recovery systems

547 Improved human and livestock waste collection and recycling systems can maintain
 548 public health while recovering freshwater, energy, and nutrients to produce more food and
 549 improving ocean health. When nutrients are recycled effectively, the food-waste-food circular
 550 economy should cost less than current systems for treating human and livestock waste that
 551 destroy nutrients, necessitating production of artificial and mined nutrients.

552 3.7.3 Solid waste resource recovery systems

553 Municipal and industrial solid waste collection systems can recover resources safely and
 554 effectively while producing energy that more than covers the cost of collection. Paying people
 555 for their solid waste would greatly reduce future marine plastic pollution. Developing countries
 556 might earn income from developed countries by exporting carbon negative biofuels.

557 3.7.4 Sustainable energy systems

558 Install multi-fuel energy systems that produce sequestration-ready CO₂. The “multi-”
 559 includes coal, natural gas, and biomass. Include ways to recycle nutrients from the energy
 560 process to grow more food and biomass-for-energy. Developing countries might earn income
 561 from developed countries by growing terrestrial biomass to fuel the developing country’s

562 electricity production and sequestering the bio-CO₂ less expensively than can be done in
 563 developed countries. Co-locate the human and solid waste resource recovery plants with
 564 sustainable energy systems for cost and circular economy synergies.

565 3.7.5 Sustainable ocean biomass-for-energy

566 Gradually scale the seafood reefs with improvements in labor productivity appropriate for
 567 satisfying global demand for liquid biofuels. Developing countries might earn income from
 568 developed countries by exporting carbon negative biofuel.

569 3.7.6 CO₂ sequestration systems

570 Employ location-appropriate CO₂ sequestration systems for the CO₂ produced and
 571 captured during energy production. Developing countries might earn income from developed
 572 countries by exporting negative carbon credits.

573 3.7.7 Floating land systems

574 Floating land (Guarino, 2019) is a collection of systems that allows people to remain in
 575 place and/or move to living on the ocean as sea levels rise.

576 3.7.8 Other public health systems

577 Both proposed paths help replace inefficient open-flame charcoal cooking with clean-
 578 burning fuel or electric stoves. They also eliminate air pollutants from electricity generation,
 579 yielding large co-benefits for air quality and human health. West et al. (2013) calculated local
 580 average marginal co-benefits of avoided mortality from air pollution ranging from \$50–
 581 380/tonne of CO₂.

582 3.7.9 All systems must be sustainable

583 In 2012 N'Yeurt et al. discussed sustainability criteria for growing macroalgae forests to
 584 reverse climate change. The technologies have evolved, and the economics have improved, now
 585 offering even more sustainability in all the ways listed by N'Yeurt et al.: environmental, climate,
 586 political, social, energy, and economic. (See SD for more discussion on how ocean forest reefs
 587 directly support twelve of the SDGs.)

588 **4 Conclusions and Recommendations**

589 4.1 Summary

590 Each country or community can pick the sustainable developments and associated
 591 technologies that best fit their resources and goals. Every sustainable development listed in
 592 Section 3.7 can start now and grow while achieving SDGs with excellent economic efficiency.
 593 These technologies can deploy to significant scale while earning profits producing seafood plus
 594 energy and nutrients from mixed plastics and organic solid waste. This is consistent with Otto et
 595 al. (2020) in that major climate efforts must be “explicitly compatible with the Sustainable
 596 Development Goals, in the sense of positive social tipping dynamics.” The health co-benefits of
 597 zero emissions energy and waste recovery strongly support this approach by generating local
 598 support, especially as these benefits are primarily local and near-term (West et al., 2013).

599 Food, jobs, adaptations for tropical oceans, and ocean SDGs are addressed while oceanic-
 600 biomass-for-energy production slowly improves. Global seafood production can increase to a
 601 billion tonnes/yr. This is nearly 6 times current seafood production (FAO, 2018) and double the
 602 combined current total meat and seafood production. (Global meat production in 2018 was 327
 603 million tonnes (Shahbandeh, 2019), with an average climate impact per kg four times that of
 604 seafood (Poore & Nemecek, 2018).)

605 The increased seafood production is supported by improved ways to ensure the nutrients
 606 people eat are recycled to land and ocean. In the ocean, recycled nutrients are distributed to
 607 macroalgae or seagrass grown on floating flexible fishing reefs positioned in the photic zone
 608 independent of seafloor depth. The fishing reefs form highly productive ecosystems supported by
 609 nutrients optimized for seasonal productivity and natural variations in nutrients and dissolved
 610 oxygen supply. Calculations suggest a billion tonnes of seafood can be grown on less than 10%
 611 of the suitable continental shelf less than 200-m seafloor depth (identified by Gentry et al.,
 612 2017). That would be about 0.3% of the world's oceans (see SS tabs 6 and 18). By growing more
 613 food in less ocean, the marine protected areas could be increased.

614 Allam Cycle power plants reduce the avoided cost (the economic penalty) to capture,
 615 compress, transport, and sequester one tonne of CO₂ from current over \$60/tonne for CCS to less
 616 than \$0/tonne for early adopters (SS tabs 14 and 16). (As “waste” sources become valuable and
 617 gases produced during Allam Cycle electricity production exceed commercial demand, the
 618 avoided cost could rise to \$26/tonne.) This significant cost drop for CCS, combined with
 619 developing country needs for new reliable electricity, provides an opportunity for developing
 620 countries to lead in healing climate change.

621 All countries can enjoy safe handling of biohazard wastes and mixed solid wastes in
 622 general with low, even negative, disposal fees using hydrothermal liquefaction (HTL) to produce
 623 22 million barrels/day (3 million tonnes/day) of biocrude oil from wastes by 2050. Additional
 624 benefits include less plastic trash reaches the ocean, less methane emissions from landfills, and
 625 profitably cleaning excess *Sargassum* off beaches.


626 4.2 Recommendations

627 The following recommendations should be implemented as soon as is feasible:

- 628 • Pause (even in the middle of construction) electricity power plant construction to check
 629 local feasibility of Allam Cycle plants. The design of Allam Cycle plants facilitates mass
 630 production. Funding agencies could purchase blocks of a thousand factory-built 300 MW
 631 power plants at a time. In addition to lower costs from mass production, this action will
 632 increase budget certainty for developing countries as they switch to Allam Cycle power.
 633 Fast start-up is encouraged while the oil industry is still buying CO₂ for enhanced oil
 634 recovery. Income from selling CO₂ for EOR will decrease the cost of electricity. (8
 635 Rivers Capital (2020) estimated in early 2020 that the global demand for CO₂ used for
 636 EOR is equivalent to nearly 6,000 of the 300 MW Allam Cycle power plants or
 637 1,800 GW.)
- 638 • Quickly build seafood-production infrastructure in developing countries.
- 639 • Convert solid waste to biocrude oil in developed and developing countries.
- 640 • Co-locate the HTL and Allam Cycle facilities to maintain the cost for capturing,
 641 compressing, and sequestering CO₂ near \$26/tonne as opposed to the \$75/tonne expense

642 for a stand-alone facility. Co-locate with other businesses and waste handling to
 643 maximize a circular economy and overall energy efficiency (e.g., pasteurizing human and
 644 medical wastes with “waste” heat and manufacturing high-performance plastics that more
 645 easily convert to biocrude or electricity).

- 646 • Take advantage of the immediate benefits from food, jobs, waste handling, and sales of
 647 liquid CO₂ byproduct to achieve SDGs. Start-up actions are further described in the SD
 648 and by Capron et al. (2020a, 2020d).
- 649 • Increase marine protected areas in proportion to the area converted to total ecosystem
 650 aquaculture.
- 651 • Establish a Legacy Carbon Fund (each country could have its own fund but with global
 652 accounting). The fund would collect more-than-the-cost of sequestering fossil-CO₂ from
 653 fossil fuel producers/users. The more-than costs are used to pay for sequestering bio-CO₂.



654
 655 Fig. 4. Global timeline to achieve SDGs and reduce atmospheric CO₂ by 2 trillion tonnes around
 656 2100 (details in SD)

657 4.3 Further Research

658 Our recommendation is to conduct research while building, operating, and maintaining
 659 commercial-scale infrastructure. The urgent needs for SDGs, addressing climate change, and
 660 assessing the effectiveness of each technology are best addressed at commercial scale. Examples
 661 of research needs include the following (see SD for additional examples):

- 662 • Life-cycle cost, planetary boundaries (Algunaibet et al., 2019) energy, and emissions
 663 analyses – The economics of reducing atmospheric CO₂ concentrations on either path
 664 appear feasible. But what are their complete life-cycle costs? Does either path exceed or
 665 increase planetary boundaries?

- 666 • Total-ecosystem aquaculture must be designed for continued biodiversity and seafood
667 production even with some fish species moving toward the poles as the tropical oceans
668 become too warm (Morley et al., 2018; Sumaila et al., 2019).
- 669 • Economics and governance – Who pays how much for achieving net-zero CO₂
670 emissions? Who pays how much for removing legacy CO₂ emissions from the
671 atmosphere for a century or so after achieving net-zero CO₂ emissions?

672 **Acknowledgments, Samples, and Data**

673 **Conflict of Interest:** The authors declare no competing interests. Authors Capron and Hasan
674 acquired knowledge of HTL processes while considering HTL manufacturers Genifuel, Algae
675 Systems, and CleanCarbon Energy for municipal biosolids and solid waste. Capron and Hasan
676 provided pro-bono advice to all three companies on topics related to recycling nutrients and
677 handling byproduct carbon (can be either fuel or food).

678 **Funding:** This research was not directly funded by agencies in the public, commercial, or not-
679 for-profit sectors. However, the authors appreciate funding by the U.S. Department of Energy's
680 Advanced Research Projects Agency - Energy (ARPA-E) MacroAlgae Research Inspiring Novel
681 Energy Resources (MARINER) Program, which helped assemble teams and facilitated work on
682 ocean macroalgae-for-energy cultivation under the following Department of Energy contracts:
683 DE-AR0000911, DE-AR0000912, DE-AR0000915, DE-AR0000916, DE-AR0000919, DE-
684 AR0000923, DE-AR0000925.

685 **Acknowledgements:**

686 Dr. Marc von Keitz for establishing and leading the U.S. Department of Energy Advanced
687 Research Projects Agency-Energy's MARINER Program.

688 Dr. Michael Huesemann and his team for providing cost and global-scale information from the
689 Pacific Northwest National Laboratory's MARINER team – Nautical Offshore Macroalgal
690 Autonomous Device (NOMAD).

691 Dr. Michael Stekoll and his team for providing cost and global-scale information from the
692 University of Alaska, Fairbanks' MARINER team – Scalable Coastal and Offshore Macroalgal
693 Farming.

694 Dr. Kelly Lucas, University of Southern Mississippi, for leading the Thad Cochran Marine
695 Aquaculture Center to propose and execute two MARINER projects: AdjustaDepth and
696 SeaweedPaddock.

697 Dr. Michael Rust of the U.S. National Oceanic and Atmospheric Administration for suggesting
698 teaming with the Thad Cochran Marine Aquaculture Center and insights concerning total
699 ecosystem aquaculture.

700 Matthew Wennerholm of AquaDam for contributing construction insights and cost data
701 informing the design, performance, and cost of containers for CO₂-hydrate storage.

702 James R. Oyler of Genifuel, Matt Atwood of Algae Systems, Craig Pichach of CleanCarbon
703 Energy, Jeffrey Moeller and Aaron Fisher of the Water Research Foundation's Leaders
704 Innovation Forum for Technology, and Greg Yamamoto of FreshMining for insights into
705 hydrothermal liquefaction, biocrude upgrading, and potential nutrient-recycling processes.

706 Rodney J. Allam, 8 Rivers, Toshiba, the U.S Department of Energy, and others associated with
 707 inventing and developing Allam Cycle electrical power plants.

708 Jade Chongsathapornpong for his pro-bono illustrations of an AdjustaDepth floating flexible reef
 709 structure with total ecosystem aquaculture while a student at Oxnard High School and
 710 edits/comments as a freshman at the Massachusetts Institute of Technology.

711 **Supplemental Document and Supplemental Spreadsheet** available online.

712 **References**

713 8 Rivers Capital. (2020). *Allam Cycle Zero Emission Coal Power*. Retrieved from
 714 <https://www.netl.doe.gov/coal/tpg/coalfirst/DirectSupercriticalCo2>

715 Alcalde, J., Flude, S., Wilkinson, M., Johnson, G., Edlmann, K., Bond, C. E., Scott, V., Gilfillan,
 716 S. M. V, Ogaya, X., & Haszeldine, R. S. (2018). Estimating geological CO₂ storage security
 717 to deliver on climate mitigation. *Nature Communications*, 9(1), 2201.
 718 <https://doi.org/10.1038/s41467-018-04423-1>

719 Aldridge, J., van der Molen, J., & Forster, R. (2012). *Wider ecological implications of*
 720 *macroalgae cultivation*. *The Crown Estate, Marine Estate Research Report*.
 721 https://www.researchgate.net/publication/285680370_Wider_ecological_implications_of_Macroalgae_cultivation

722

723 Algunaibet, I. M., Pozo, C., Galán-Martín, Á., Huijbregts, M. A. J., Mac Dowell, N., & Guillén-
 724 Gosálbez, G. (2019). Powering sustainable development within planetary boundaries. *Energy*
 725 *& Environmental Science*, 12(6), 1890–1900. <https://doi.org/10.1039/C8EE03423K>

726 Allam, R., Martin, S., Forrest, B., Fetvedt, J., Lu, X., Freed, D., Brown, G. W., Sasaki, T., Itoh,
 727 M., & Manning, J. (2017). Demonstration of the Allam Cycle: An Update on the
 728 Development Status of a High Efficiency Supercritical Carbon Dioxide Power Process
 729 Employing Full Carbon Capture. *Energy Procedia*, 114(November 2016), 5948–5966.
 730 <https://doi.org/10.1016/j.egypro.2017.03.1731>

731 Atlas-Copco. (2020). *CO₂ Compressors*. Retrieved from <https://www.atlascopco.com/en-us/compressors/products/gas-compressors/co2-compressor>

732

733 Baker, S. E., Stolaroff, J. K., Peridas, G., Pang, S. H., Goldstein, H. M., Lucci, F. R., et al.
 734 (2020). *Getting to Neutral: Options for Negative Carbon Emissions in California*. The
 735 ASHA Leader. Retrieved from <https://www.llnl.gov/news/new-lab-report-outlines-ways-california-could-reach-goal-becoming-carbon-neutral-2045>

736

737 Boerner, L. K. (2019). Industrial ammonia production emits more CO₂ than any other chemical-
 738 making reaction. Chemists want to change that. *Chemical & Engineering News*, 97 (24).
 739 <https://cen.acs.org/environment/green-chemistry/Industrial-ammonia-production-emits-CO2/97/i24>

740

741 Boyd, P.W., Sundby, S., and Pörtner, H.-O. (2014). Cross-chapter box on Net Primary
 742 Production in the Ocean. In Field, C.B., V.R. Barros, D.J. Dokken, K.J. Mach, M.D.
 743 Mastrandrea, T.E. Bilir, M. Chatterjee, K.L. Ebi, Y.O. Estrada, R.C. Genova, B. Girma, E.S.
 744 Kissel, A.N. Levy, S. MacCracken, P.R. Mastrandrea (Ed.), *Climate Change 2014: Impacts,*
 745 *Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working*
 746 *Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*

- 747 (pp. 133–136). Cambridge University Press. Retrieved from
 748 <https://www.ipcc.ch/report/ar5/wg2/>
- 749 BP. (2019). *BP Statistical Review of World Energy*. Retrieved from
 750 [https://www.bp.com/en/global/corporate/energy-economics/statistical-review-of-world-](https://www.bp.com/en/global/corporate/energy-economics/statistical-review-of-world-energy.html)
 751 [energy.html](https://www.bp.com/en/global/corporate/energy-economics/statistical-review-of-world-energy.html)
- 752 Bradshaw, J., & Dance, T. (2005). Mapping geological storage prospectivity of CO₂ for the
 753 world's sedimentary basins and regional source to sink matching. *Greenhouse Gas Control*
 754 *Technologies, August*, 583–591. <https://doi.org/10.1016/B978-008044704-9/50059-8>
- 755 Brewer, P. G. (2000). Roger Revelle commemorative lecture contemplating action: Storing
 756 carbon dioxide in the ocean. *Oceanography, 13*(SPL.ISS. 2), 84–92.
 757 <https://doi.org/10.5670/oceanog.2000.38>
- 758 Brown, T. (2016). *Ammonia production causes 1% of total global GHG emissions*. Retrieved
 759 from [https://ammoniaindustry.com/ammonia-production-causes-1-percent-of-total-global-](https://ammoniaindustry.com/ammonia-production-causes-1-percent-of-total-global-ghg-emissions/)
 760 [ghg-emissions/](https://ammoniaindustry.com/ammonia-production-causes-1-percent-of-total-global-ghg-emissions/)
- 761 Bruun, J. (2019). *Revolutionizing the way we manage waste*. Retrieved from
 762 [https://eng.au.dk/en/news-and-events/news/show/artikel/revolutionising-the-way-we-](https://eng.au.dk/en/news-and-events/news/show/artikel/revolutionising-the-way-we-manage-waste-a-danish-researcher-is-developing-a-pioneering-new-technolo-1/)
 763 [manage-waste-a-danish-researcher-is-developing-a-pioneering-new-technolo-1/](https://eng.au.dk/en/news-and-events/news/show/artikel/revolutionising-the-way-we-manage-waste-a-danish-researcher-is-developing-a-pioneering-new-technolo-1/)
- 764 Buschmann, A. H., Camus, C., Infante, J., Neori, A., Israel, Á., Hernández-González, M. C.,
 765 Pereda, S. V, Gomez-Pinchetti, J. L., Golberg, A., Tadmor-Shalev, N., & Critchley, A. T.
 766 (2017). Seaweed production: overview of the global state of exploitation, farming and
 767 emerging research activity. *European Journal of Phycology, 52*(4), 391–406.
 768 <https://doi.org/10.1080/09670262.2017.1365175>
- 769 California Air Resources Board. (2020). *California-Quebec Joint Auction Settlement Prices*.
 770 Retrieved from https://ww3.arb.ca.gov/cc/capandtrade/auction/results_summary.pdf
- 771 CalRecycle. (2014). *Disposal-Facility-Based Characterization of Solid Waste in California*.
 772 *Publication # DRRR-2015-1546*. Retrieved from
 773 <https://www2.calrecycle.ca.gov/Publications/Details/1546>
- 774 Capo, T. R., Jaramillo, J. C., Boyd, A. E., Lapointe, B. E., & Serafy, J. E. (1999). Sustained high
 775 yields of *Gracilaria* (Rhodophyta) grown in intensive large-scale culture, *Journal of Applied*
 776 *Phycology, 11*(2), 143. <https://doi.org/10.1023/A:1008077722769>
- 777 Capron, M. E. (2019). *Centuries of food and job security for coastal peoples*. Contest Entry in
 778 BridgeBuilder™ 2019 Challenge: People on the Move. Retrieved from
 779 [https://challenges.openideo.com/challenge/2019-bridgebuilder-challenge/ideas/centuries-of-](https://challenges.openideo.com/challenge/2019-bridgebuilder-challenge/ideas/centuries-of-food-and-job-security-for-coastal-peoples)
 780 [food-and-job-security-for-coastal-peoples](https://challenges.openideo.com/challenge/2019-bridgebuilder-challenge/ideas/centuries-of-food-and-job-security-for-coastal-peoples)
- 781 Capron, M. E., & Piper, D. (2019). *Abundant Food and Export Income with Total ecosystem*
 782 *Aquaculture*. Contest Entry in Reshaping Development Pathways in LDCs. Retrieved from
 783 [https://www.climatecolab.org/contests/2019/reshapingdevelopmentpathwaysinLDCs/c/propo-](https://www.climatecolab.org/contests/2019/reshapingdevelopmentpathwaysinLDCs/c/proposal/1334620)
 784 [sal/1334620](https://www.climatecolab.org/contests/2019/reshapingdevelopmentpathwaysinLDCs/c/proposal/1334620)
- 785 Capron, M. E., Lucas, K, Chambers, MD, Lapointe, B, N'Yeurt, A. de R., Krueger-Hadfield, SA,
 786 et al. (2019). Building New Fishery Ecosystems: The Science of Flexible Floating Fishing
 787 Reefs. *International Symposium on Stock Enhancement & Sea Ranching*. Retrieved from
 788 <https://mote.org/media/uploads/contact/ISSESR-Wednesday-1150-Piper.pdf>

- 789 Capron, M. E., Lucas, K., Chambers, M. D., & Stewart, J. R. (2018). Ocean Forests :
 790 Breakthrough Yields for Macroalgae. *OCEANS 2018 MTS/IEEE Charleston*, 1–6.
 791 <https://doi.org/10.1109/OCEANS.2018.8604586>
- 792 Capron, M. E., N'Yeurt, A. de R., Kim, J. K., Pichach, C., Chambers, M. D., Fuhrman, R., et al.
 793 (2020a). Reversing Climate Change within 100 years: The scale to restore natural CO₂ levels.
 794 *Ocean Sciences Meeting Presentation SI41A-02*. Retrieved from
 795 <https://agu.confex.com/agu/osm20/meetingapp.cgi/Paper/648119>
- 796 Capron, M. E., Prasad, R., N'Yeurt, A. de R., Hewavitharane, C., & Stewart, J. R. (2020b). *Total*
 797 *Ecosystem Aquaculture - Sustainable Seafood Production for Fiji, the Pacific, and then the*
 798 *World*. Contest Entry in Rockefeller Foundation Food System Vision Prize. Retrieved from
 799 [https://challenges.openideo.com/challenge/food-system-vision-prize/open-](https://challenges.openideo.com/challenge/food-system-vision-prize/open-submission/restorative-aquaculture-sustainable-seafood-production-for-the-world)
 800 [submission/restorative-aquaculture-sustainable-seafood-production-for-the-world](https://challenges.openideo.com/challenge/food-system-vision-prize/open-submission/restorative-aquaculture-sustainable-seafood-production-for-the-world)
- 801 Capron, M. E., Prasad, R., N'Yeurt, A. de R., Hopkins, K., Stewart, J. R., Hasan, M. A., et al.
 802 (2020c). Increasing South Pacific aquaculture with complete ecosystems. *Ocean Sciences*
 803 *Meeting Presentation SI12A-05*. Retrieved from
 804 <https://agu.confex.com/agu/osm20/meetingapp.cgi/Paper/649244>
- 805 Capron, M. E., Stewart, J. R., & Rowe, R. (2013). Secure Seafloor Container CO₂ Storage.
 806 *OCEANS'13 MTS/IEEE San Diego Technical Program #130503-115*.
 807 <https://doi.org/10.23919/OCEANS.2013.6741182>
- 808 Capron, M. E., Stewart, J. R., N'Yeurt, A. de R., Prasad, R., Hewavitharane, C., Hasan, M. A., et
 809 al. (2020d). *Sustainable Development that supports Ocean Science. Ocean Science that*
 810 *supports Sustainable Development. Concepts for the UN Decade of Ocean Science for*
 811 *Sustainable Development*. Retrieved from [https://oceandecade.org/resource/102/Planning-](https://oceandecade.org/resource/102/Planning-Concrete-Deliverables-for-the-UN-Decade-of-Ocean-Sciences-for-Sustainable-Development-2021-2030)
 812 [Concrete-Deliverables-for-the-UN-Decade-of-Ocean-Sciences-for-Sustainable-](https://oceandecade.org/resource/102/Planning-Concrete-Deliverables-for-the-UN-Decade-of-Ocean-Sciences-for-Sustainable-Development-2021-2030)
 813 [Development-2021-2030](https://oceandecade.org/resource/102/Planning-Concrete-Deliverables-for-the-UN-Decade-of-Ocean-Sciences-for-Sustainable-Development-2021-2030)
- 814 Chambers, M. D. (2013). *Integrated Multi-Trophic Aquaculture of Steelhead, Blue Mussels and*
 815 *Sugar Kelp*. PhD dissertation, University of New Hampshire, Durham, NH.
- 816 Chan, F., Barth, J. A., Blanchette, C. A., Byrne, R. H., Chavez, F., Cheriton, O., Feely, R. A.,
 817 Friederich, G., Gaylord, B., Gouhier, T., Hacker, S., Hill, T., Hofmann, G., McManus, M. A.,
 818 Menge, B. A., Nielsen, K. J., Russell, A., Sanford, E., Sevdjian, J., & Washburn, L. (2017).
 819 Persistent spatial structuring of coastal ocean acidification in the California Current System.
 820 *Scientific Reports*, 7(1), 2526. <https://doi.org/10.1038/s41598-017-02777-y>
- 821 Chen, S. (2018). *The development of sea surface carbonate chemistry in the coastal upwelling*
 822 *area off Peru following a simulated OMZ upwelling event*. Thesis submitted to Kiel
 823 University, Kiel, Germany.
- 824 Choi, Y.-S., Young, D., Nešić, S., & Gray, L. G. S. (2013). Wellbore integrity and corrosion of
 825 carbon steel in CO₂ geologic storage environments: A literature review. *International*
 826 *Journal of Greenhouse Gas Control*, 16, S70–S77.
 827 <https://doi.org/10.1016/J.IJGGC.2012.12.028>
- 828 Chung, I. K., Oak, J. H., Lee, J. A., Shin, J. A., Kim, J. G., & Park, K.-S. (2013). adaptation
 829 against global warming : Korean Project Overview. *ICES Journal of Marine Science*,
 830 68(November), 66–74. <https://doi.org/10.1093/icesjms/fss206>

- 831 Clarke, S. & Preto, F. (2011). Biomass Burn Characteristics. Ministry of Agriculture, Food and
 832 Rural Affairs, Ontario, Canada. Retrieved from
 833 <http://www.omafra.gov.on.ca/english/engineer/facts/11-033.pdf>
- 834 Cottier-Cook, E. J., Nagabhushan, N., Badis, Y., Campbell, M. L., Chopin, T., Fang, J., ... Jiang,
 835 Z. (2016). Safeguarding the future of the global seaweed aquaculture industry. United
 836 Nations University (INWEH) and Scottish Association for Marine Science Policy Brief.
 837 ISBN 978-92-808-6080-1. Retrieved from [http://inweh.unu.edu/safeguarding-the-future-of-](http://inweh.unu.edu/safeguarding-the-future-of-the-global-seaweed-aquaculture-industry-policy-brief/)
 838 [the-global-seaweed-aquaculture-industry-policy-brief/](http://inweh.unu.edu/safeguarding-the-future-of-the-global-seaweed-aquaculture-industry-policy-brief/)
- 839 D'Angelo, C. (2010). Historic California Kelp Harvest, 1916 – 2009. National Marine Fisheries
 840 Service report. Retrieved from
 841 [https://drive.google.com/file/d/1hVQXdt2TrdVB2LRpnWp1SGvs7XqDE3re/view?usp=shar](https://drive.google.com/file/d/1hVQXdt2TrdVB2LRpnWp1SGvs7XqDE3re/view?usp=sharing)
 842 [ing](https://drive.google.com/file/d/1hVQXdt2TrdVB2LRpnWp1SGvs7XqDE3re/view?usp=sharing)
- 843 Daioglou, V, Stehfest, E, Wicke, B, Faaij, A. (2016). Projections-of-the-availability-and-cost-of-
 844 residues-from-agriculture-and-forestry.pdf. *Bioenergy*, 8, 456–470.
 845 <https://doi.org/10.1111/gcbb.12285>
- 846 Daly, C. and Halbleib, M. (2014). *Potential Yield Mapping of Dedicated Energy Crops*.
 847 Retrieved from http://energy.gov/sites/prod/files/2014/11/f19/daly_biomass_2014.pdf
- 848 Das, PK, Das, BP, Dash, P. (2019). Role of Energy Crops to Meet the Rural Energy Needs: An
 849 Overview. In *Biomass Valorization to Bioenergy*. [https://doi.org/10.1007/978-981-15-0410-](https://doi.org/10.1007/978-981-15-0410-5_2)
 850 [5_2](https://doi.org/10.1007/978-981-15-0410-5_2)
- 851 Deng, H., Bielicki, J. M., Oppenheimer, M., Fitts, J. P., & Peters, C. A. (2017). Leakage risks of
 852 geologic CO₂ storage and the impacts on the global energy system and climate change
 853 mitigation. *Climatic Change*, 144(2), 151–163. <https://doi.org/10.1007/s10584-017-2035-8>
- 854 Deng, Y. Y., Koper, M., Haigh, M., & Dornburg, V. (2015). Country-level assessment of long-
 855 term global bioenergy potential. *Biomass and Bioenergy*, 74, 253–267.
 856 <https://doi.org/https://doi.org/10.1016/j.biombioe.2014.12.003>
- 857 Devi, S., Gupta, C., Jat, S. L. & Parmar, M. S. (2017). Crop residue recycling for economic and
 858 environmental sustainability: The case of India. *Open Agriculture* 2, 486–494.
 859 <https://doi.org/10.1515/opag-2017-0053>
- 860 Duarte, C. M. (1992). Nutrient concentration of aquatic plants: Patterns across species.
 861 *Limnology and Oceanography*, 37(4), 882–889. <https://doi.org/10.4319/lo.1992.37.4.0882>
- 862 Eisentraut, A. (2010). *Sustainable production of second-generation biofuels: potential and*
 863 *perspectives in major economies and developing countries*.
 864 <https://doi.org/10.1787/5kmh3njpt6r0-en>
- 865 Enerdata. (2019). *Global Energy Statistical Yearbook*. Retrieved from
 866 <https://yearbook.enerdata.net/electricity/electricity-domestic-consumption-data.html>
- 867 Environmental Research & Education Foundation. (2019). *Analysis of MSW Tipping Fees*.
 868 Retrieved from <http://www.erefnd.org>
- 869 Eriksson, O. & Finnveden, G. (2009). Plastic waste as a fuel – CO₂-neutral or not? *Energy and*
 870 *Environmental Science*, 2, 907–914. DOI: 10.1039/b908135f

- 871 Faaij, A. (Lead Author) (2007). Potential Contribution of Bioenergy to the World's Future
 872 Energy Demand, IEA Bioenergy Strategic Position Paper, IEA Bioenergy:ExCo:2007:02.
 873 Retrieved from [https://www.ieabioenergy.com/publications/potential-contribution-of-](https://www.ieabioenergy.com/publications/potential-contribution-of-bioenergy-to-the-worlds-future-energy-demand)
 874 [bioenergy-to-the-worlds-future-energy-demand](https://www.ieabioenergy.com/publications/potential-contribution-of-bioenergy-to-the-worlds-future-energy-demand)
- 875 FAO. (2018). *The State of World Fisheries and Aquaculture 2018 - Meeting the sustainable*
 876 *development goals*. Retrieved from <http://www.fao.org/documents/card/en/c/I9540EN/>
- 877 FAO. (2015). *World fertilizer trends and outlook to 2018*. ISBN 978-92-5-108692-6. Retrieved
 878 from <http://www.fao.org/3/a-i4324e.pdf>
- 879 Feely, R. A., Sabine, C. L., Hernandez-Ayon, J. M., Ianson, D., & Hales, B. (2008). Evidence for
 880 Upwelling of Corrosive “Acidified” Water onto the Continental Shelf. *Science*, *320*(5882),
 881 1490–1492. <https://doi.org/10.1126/science.1155676>
- 882 Fernandes, D., Wang, S., Xu, Q., Buss, R., & Chen, D. (2019). Process and Carbon Footprint
 883 Analyses of the Allam Cycle Power Plant Integrated with an Air Separation Unit. *Clean*
 884 *Technologies*, *1*(1), 325–340. <https://doi.org/10.3390/cleantechnol1010022>
- 885 Field, C. B., Behrenfeld, M. J., Randerson, J. T., & Falkowski, P. (1998). Primary Production of
 886 the Biosphere: Integrating Terrestrial and Oceanic Components. *Science*, *281*(5374), 237 LP
 887 – 240. <https://doi.org/10.1126/science.281.5374.237>
- 888 Fischer, G. & Schrattenholzer, L. (2001), “Global Bioenergy Potentials Through 2050,” *Biomass*
 889 *and Bioenergy*, Vol. 20, pp. 151-159. [https://doi.org/10.1016/S0961-9534\(00\)00074-X](https://doi.org/10.1016/S0961-9534(00)00074-X)
- 890 Flannery, T. (2015). *Atmosphere of Hope: Searching for Solutions to the Climate Crisis*. Boston:
 891 Atlantic Monthly Press.
- 892 Flannery, T. (2017). *Sunlight and Seaweed, an Argument for How to Feed, Power and Clean Up*
 893 *the World*. Melbourne, Australia: The Text Publishing Company.
- 894 Frank, A. B., Berdahl, J. D., Hanson, J. D., Liebig, M. A., & Johnson, H. A. (2004). Biomass and
 895 Carbon Partitioning in Switchgrass. *Crop Science*, *44*, 1391–1396.
 896 <https://doi.org/10.2135/cropsci2004.1391>
- 897 Friedlingstein, P., Jones, M. W., O’Sullivan, M., Andrew, R. M., Hauck, J., Peters, G. P., et al.
 898 (2019). Global Carbon Budget 2019. *Earth System Science Data*, *11*(4), 1783–1838.
 899 <https://doi.org/10.5194/essd-11-1783-2019>
- 900 Fuss, S., Lamb, W. F., Callaghan, M. W., Hilaire, J., Creutzig, F., Amann, T., et al. (2018).
 901 Negative emissions - Part 2: Costs, potentials and side effects. *Environmental Research*
 902 *Letters*, *13*(6). <https://doi.org/10.1088/1748-9326/aabf9f>
- 903 Future Timeline. (2018). World GDP per capita, 1950-2100. Retrieved from
 904 <https://www.futuretimeline.net/data-trends/8.htm>
- 905 Genifuel. (2019). *Clean Conversion of Wet Waste to Fuel*. Retrieved from
 906 <https://www.genifuel.com/>
- 907 Gentry, R. R., Froehlich, H. E., Grimm, D., Kareiva, P., Parke, M., Rust, M., Gaines, S. D., &
 908 Halpern, B. S. (2017). Mapping the global potential for marine aquaculture. *Nature Ecology*
 909 *& Evolution*, *1*(9), 1317–1324. <https://doi.org/10.1038/s41559-017-0257-9>
- 910 GESAMP. (2019). High level review of a wide range of proposed marine geoengineering
 911 techniques. Boyd, P.W. and Vivian, C.M.G. eds. (IMO/FAO/UNESCO-

- 912 IOC/UNIDO/WMO/IAEA/UN/UN Environment/ UNDP/ISA Joint Group of Experts on the
 913 Scientific Aspects of Marine Environmental Protection). GESAMP Report No. 98. Retrieved
 914 from [http://www.gesamp.org/publications/high-level-review-of-a-wide-range-of-proposed-](http://www.gesamp.org/publications/high-level-review-of-a-wide-range-of-proposed-marine-geoengineering-techniques)
 915 [marine-geoengineering-techniques](http://www.gesamp.org/publications/high-level-review-of-a-wide-range-of-proposed-marine-geoengineering-techniques)
- 916 Gíslason, S. R., Sigurdardóttir, H., Aradóttir, E. S., & Oelkers, E. H. (2018). A brief history of
 917 CarbFix: Challenges and victories of the project's pilot phase. *Energy Procedia*, *146*, 103–
 918 114. <https://doi.org/10.1016/j.egypro.2018.07.014>
- 919 Global CCS Institute. (2019). *CO₂RE Database*. Retrieved from <https://co2re.co>
- 920 Gough, C. (2011). *Terrestrial Primary Production: Fuel for Life*. Retrieved from
 921 [https://www.nature.com/scitable/knowledge/library/terrestrial-primary-production-fuel-for-](https://www.nature.com/scitable/knowledge/library/terrestrial-primary-production-fuel-for-life-17567411/)
 922 [life-17567411/](https://www.nature.com/scitable/knowledge/library/terrestrial-primary-production-fuel-for-life-17567411/)
- 923 Guarino, B. (2019). As seas rise, the U.N. explores a bold plan: Floating cities. *Washington Post*.
 924 Retrieved from [https://www.washingtonpost.com/science/2019/04/05/seas-rise-un-explores-](https://www.washingtonpost.com/science/2019/04/05/seas-rise-un-explores-bold-plan-floating-cities/)
 925 [bold-plan-floating-cities/](https://www.washingtonpost.com/science/2019/04/05/seas-rise-un-explores-bold-plan-floating-cities/)
- 926 Gunnarsson, I., Aradóttir, E. S., Oelkers, E. H., Clark, D. E., Arnarson, M. P., Sigfússon, B., et
 927 al. (2018). The rapid and cost-effective capture and subsurface mineral storage of carbon and
 928 sulfur at the CarbFix2 site. *International Journal of Greenhouse Gas Control*, *79*, 117–126.
 929 <https://doi.org/10.1016/J.IJGGC.2018.08.014>
- 930 Hall, C. A. S., Dale, B. E., & Pimentel, D. (2011). Seeking to understand the reasons for
 931 different energy return on investment (EROI) estimates for biofuels. *Sustainability*, *3*(12),
 932 2413–2432. <https://doi.org/10.3390/su3122413>
- 933 Hartmann, J. & Moosdorf, N. (2012). The new global lithological map database GLiM: a
 934 representation of rock properties at the Earth surface. *Geochem. Geophys. Geosyst.* **13**,
 935 Q12004. <https://doi.org/10.1029/2012GC004370>
- 936 Hastings, R. A., Rutterford, L. A., Freer, J. J., Collins, R. A., Simpson, S. D., & Genner, M. J.
 937 (2020). Climate Change Drives Poleward Increases and Equatorward Declines in Marine
 938 Species. *Current Biology*, *30*(8), 1572-1577.e2.
 939 <https://doi.org/https://doi.org/10.1016/j.cub.2020.02.043>
- 940 Hepburn, C., Adlen, E., Beddington, J., Carter, E. A., Fuss, S., Mac Dowell, N., et al. (2019).
 941 The technological and economic prospects for CO₂ utilization and removal. *Nature*,
 942 *575*(7781), 87–97. <https://doi.org/10.1038/s41586-019-1681-6>
- 943 Hicks, C. C., Cohen, P. J., Graham, N. A. J., Nash, K. L., Allison, E. H., D'Lima, C., et al. 2019.
 944 Harnessing global fisheries to tackle micronutrient deficiencies. *Nature* **574**, 95–98.
 945 [doi:10.1038/s41586-019-1592-6](https://doi.org/10.1038/s41586-019-1592-6)
- 946 Hoogwijk, M., Faaij, A., de Vries, B., Turkenburg, W. C. (2009). Exploration of regional and
 947 global cost-supply curves of biomass from short-rotation crops at abandoned cropland and
 948 rest land under four IPCC SRES land-use scenarios. *Biomass and Bioenergy* *33*, 26– 43.
 949 [doi:10.1016/j.biombioe.2008.04.005](https://doi.org/10.1016/j.biombioe.2008.04.005)
- 950 Hudiburg, T. W., Luysaert, S., Thornton, P. E., & Law, B. E. (2013). Interactive Effects of
 951 Environmental Change and Management Strategies on Regional Forest Carbon Emissions.

- 952 *Environmental Science & Technology*, 47(22), 13132–13140.
 953 <https://doi.org/10.1021/es402903u>
- 954 Huesemann, M., Mumford, T., Wood, G., Quinn, J., Wang, T., Yang, Z. (2017). *Nautical*
 955 *Offshore Macroalgal Autonomous Device (NOMAD)*. Retrieved from [https://arpa-](https://arpa-e.energy.gov/?q=slick-sheet-project/nautical-offshore-macroalgal-autonomous-device)
 956 [e.energy.gov/?q=slick-sheet-project/nautical-offshore-macroalgal-autonomous-device](https://arpa-e.energy.gov/?q=slick-sheet-project/nautical-offshore-macroalgal-autonomous-device)
- 957 Ibisworld. (2019). *Global Coal Mining Industry - Market Research Report*. Retrieved from
 958 <https://www.ibisworld.com/global/market-research-reports/global-coal-mining-industry>
- 959 IEA. (2018). *World Energy Outlook 2018*. Retrieved from [https://www.iea.org/reports/world-](https://www.iea.org/reports/world-energy-outlook-2018)
 960 [energy-outlook-2018](https://www.iea.org/reports/world-energy-outlook-2018)
- 961 Investopedia. (2020). *What percentage of the global economy consists of the oil and gas drilling*
 962 *sector?* Retrieved from [https://www.investopedia.com/ask/answers/030915/what-percentage-](https://www.investopedia.com/ask/answers/030915/what-percentage-global-economy-comprised-oil-gas-drilling-sector.asp)
 963 [global-economy-comprised-oil-gas-drilling-sector.asp](https://www.investopedia.com/ask/answers/030915/what-percentage-global-economy-comprised-oil-gas-drilling-sector.asp)
- 964 Ioelovich, M. (2015). Recent Findings and the Energetic Potential of Plant Biomass as a
 965 Renewable Source of Biofuels – A Review. *BioResources*, 10(1), 1879-1914. Retrieved from
 966 [https://ojs.cnr.ncsu.edu/index.php/BioRes/article/view/BioRes_10_1_Ioelovich_Review_Pla](https://ojs.cnr.ncsu.edu/index.php/BioRes/article/view/BioRes_10_1_Ioelovich_Review_Plant_Biomass_Renewable_Source_Biofuels)
 967 [nt_Biomass_Renewable_Source_Biofuels](https://ojs.cnr.ncsu.edu/index.php/BioRes/article/view/BioRes_10_1_Ioelovich_Review_Plant_Biomass_Renewable_Source_Biofuels)
- 968 IPCC. (2014). *Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II*
 969 *and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*
 970 *[R.K. Pachauri and L.A. Meyer (eds.)]*. Retrieved from <https://www.ipcc.ch/report/ar5/syr/>
- 971 IPCC. (2018). *Global Warming of 1.5°C. An IPCC Special Report on the impacts of global*
 972 *warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission*
 973 *pathways, in the context of strengthening the global response to the threat of climate change,*
 974 *sustainable development, and efforts to eradicate poverty*. Retrieved from
 975 <https://www.ipcc.ch/sr15/download/>
- 976 Irlam, L. (2017). Global costs of carbon capture and storage. Report for the Global CCS Institute.
 977 Retrieved from [https://www.globalccsinstitute.com/archive/hub/publications/201688/global-](https://www.globalccsinstitute.com/archive/hub/publications/201688/global-ccs-cost-updatev4.pdf)
 978 [ccs-cost-updatev4.pdf](https://www.globalccsinstitute.com/archive/hub/publications/201688/global-ccs-cost-updatev4.pdf)
- 979 Jacobson, M. Z. (2019). The Health and Climate Impacts of Carbon Capture and Direct Air
 980 Capture, *Energy & Environmental Science*. <https://doi.org/10.1039/C9EE02709B>
- 981 Jacobson, M. Z. (2020). Why Not Synthetic Direct Air Carbon Capture and Storage (SDACCS)
 982 as Part of a 100% Wind-Water-Solar (WWS) and Storage Solution to Global Warming, Air
 983 Pollution, and Energy Security. In *100% Clean, Renewable Energy and Storage for*
 984 *Everything*, Textbook in press, *Cambridge University Press*. Retrieved from
 985 <https://web.stanford.edu/group/efmh/jacobson/WWSBook/WWSBook.html>
- 986 James, R., Kearins, D., Turner, M., Woods, M., Kuehn, N., Zoelle, A. (2019). Cost and
 987 Performance Baseline for Fossil Energy Plants Volume 1: Bituminous Coal and Natural Gas
 988 to Electricity. Report prepared by MESA for the U.S. DOE NETL. DOI: 10.2172/1569246
- 989 Jiang, Y., Jones, S. B., Zhu, Y., Snowden-Swan, L., Schmidt, A. J., Billing, J. M., & Anderson,
 990 D. (2019). Techno-economic uncertainty quantification of algal-derived biocrude via
 991 hydrothermal liquefaction. *Algal Research*, 39, 101450.
 992 <https://doi.org/10.1016/J.ALGAL.2019.101450>

- 993 Jiao, J. P., Wang, F., Duan, P. G., Xu, Y. P., Yan, W. H. (2017). Catalytic hydrothermal
 994 gasification of microalgae for producing hydrogen and methane-rich gas *Energy Sources,*
 995 Part A: Recovery, Utilization, and Environmental Effects Volume 39, 2017 – Issue 9.
 996 <https://doi.org/10.1080/15567036.2016.1270375>
- 997 Johansson, L., Zahirovic, S. & Müller, R. D. (2018). The interplay between the eruption and
 998 weathering of Large Igneous Provinces and the deep-time carbon cycle. *Geophys. Res. Lett.*
 999 **45**, 5380–5389. <https://doi.org/10.1029/2017GL076691>
- 1000 Jolly, R. (2004). Dutch Industrial waste continues to decrease. Retrieved from
 1001 <https://www.cbs.nl/en-gb/news/2004/42/industrial-waste-continues-to-decrease>
- 1002 Kaza, S., Yao, L., Bhada-Tata, P., and Van Woerden, F. (2018). What a Waste 2.0: A Global
 1003 Snapshot of Solid Waste Management to 2050. The World Bank.
 1004 <https://doi.org/doi:10.1596/978-1-4648-1329-0>
- 1005 Kelemen, P. (1998). Melt extraction from the mantle beneath mid-ocean ridges. *Oceanus*, *41*,
 1006 23–28. Retrieved from [https://www.whoi.edu/oceanus/feature/melt-extraction-from-the-](https://www.whoi.edu/oceanus/feature/melt-extraction-from-the-mantle-beneath-mid-ocean-ridges/)
 1007 [mantle-beneath-mid-ocean-ridges/](https://www.whoi.edu/oceanus/feature/melt-extraction-from-the-mantle-beneath-mid-ocean-ridges/)
- 1008 Kelemen, P., Benson, S. M., Pilorgé, H., Psarras, P., & Wilcox, J. (2019). An Overview of the
 1009 Status and Challenges of CO₂ Storage in Minerals and Geological Formations. *Frontiers in*
 1010 *Climate*, *1*, 9. <https://doi.org/10.3389/fclim.2019.00009>
- 1011 Keller, D. P., Lenton, A., Littleton, E. W., Oschlies, A., Scott, V., & Vaughan, N. E. (2018). The
 1012 Effects of Carbon Dioxide Removal on the Carbon Cycle. *Current Climate Change Reports*,
 1013 *4*(3), 250–265. <https://doi.org/10.1007/s40641-018-0104-3>
- 1014 Kim, J. K., Yarish, C., Hwang, E. K., Park, M., & Kim, Y. (2017). Seaweed aquaculture:
 1015 cultivation technologies, challenges and its ecosystem services. *ALGAE*, *32*(1), 1–13.
 1016 <https://doi.org/10.4490/algae.2017.32.3.3>
- 1017 Kim, J. K., Wu, H., Shin, S. K., Park, M. S., Kim, Y. D., Do, Y. H., Kim, N. G., Han, T. (2017).
 1018 An effective integrated multi-trophic aquaculture system for Korean waters. In *The 2nd*
 1019 *International Integrated Multi-Trophic Aquaculture Symposium Book of Abstracts* (pp. 17–
 1020 32).
- 1021 Knowler, D., Chopin, T., Martínez-Espiñeira, R., Neori, A., Nobre, A., Noce, A., & Reid, G.
 1022 (2020). The economics of Integrated Multi-Trophic Aquaculture: where are we now and
 1023 where do we need to go? *Reviews in Aquaculture*. <https://doi.org/10.1111/raq.12399>
- 1024 Knutti, R., & Rogelj, J. (2015). The legacy of our CO₂ emissions: a clash of scientific facts,
 1025 politics and ethics. *Climatic Change*, *133*(3), 361–373. [https://doi.org/10.1007/s10584-015-](https://doi.org/10.1007/s10584-015-1340-3)
 1026 [1340-3](https://doi.org/10.1007/s10584-015-1340-3)
- 1027 Köhn, E. E., Thomsen, S., Arévalo-Martinez, D. L., & Kanzow, T. (2017). Submesoscale CO₂
 1028 variability across an upwelling front off Peru. *Ocean Science*, *13*(6), 1017–1033.
 1029 <https://doi.org/10.5194/os-13-1017-2017>
- 1030 Korai, M. S., Mahar, R. B., Uqaili, M. A. (2014). Assessment of Power Generation Potential
 1031 from Municipal Solid Wastes: A Case Study of Hyderabad City, Sindh, Pakistan. *Pakistan*
 1032 *Journal of Analytical and Environmental Chemistry* *15* (1) 18 – 27.
 1033 <https://doaj.org/article/cf1958f16a2143f2829a43b64a4540de>

- 1034 Kostecki, J., Greinert, A., Drab, M., Wasylewicz, R., Szafraniec, M., Stodulski, G., Wypych, M.
 1035 (2014). The total content of nitrogen in leaves and wood of trees growing in the area affected
 1036 by the Glogow Copper Smelter. *Journal of Elementology* 1:137-148.
 1037 <https://doi.org/10.5601/jelem.2014.19.4.401>
- 1038 Krause-Jensen, D., Lavery, P., Serrano, O., Marba, N., Masque, P., & Duarte, C. M. (2018).
 1039 Sequestration of macroalgal carbon: The elephant in the Blue Carbon room. *Biology Letters*,
 1040 14(6). <https://doi.org/10.1098/rsbl.2018.0236>
- 1041 Krevor, S. C., Graves, C. R., Van Gosen, B. S., & McCafferty, A. E. (2009). Mapping the
 1042 mineral resource base for mineral carbon-dioxide sequestration in the conterminous United
 1043 States. In *Data Series* (Version 1). <https://doi.org/10.3133/ds414>
- 1044 Krueger-Hadfield, S. A. & Hoban, S. M. (2015). The importance of effective sampling for
 1045 exploring the population dynamics of haploid-diploid seaweeds. *Journal of Phycology* 52:1–
 1046 9. <https://doi.org/10.1111/jpy.12366>
- 1047 Kumar, P., Kumar, S., & Joshi, L. (2015). *Socioeconomic and Environmental Implications of*
 1048 *Agricultural Residue Burning: A Case Study of Punjab, India*. Springer: New Delhi
 1049 Heidelberg New York Dordrecht London.
- 1050 Kunzig, R. (2020). The End of Trash. *National Geographic* March 2020. Retrieved from
 1051 <https://www.nationalgeographic.com/magazine/2020/03/>
- 1052 Lapointe, B. E. (1985), Strategies for pulsed nutrient supply to *Gracilaria* cultures in the Florida
 1053 Keys: Interactions between concentration and frequency of nutrient pulses, *Journal of*
 1054 *Experimental Marine Biology and Ecology*, 3, 211-222. [https://doi.org/10.1016/0022-](https://doi.org/10.1016/0022-0981(85)90240-0)
 1055 [0981\(85\)90240-0](https://doi.org/10.1016/0022-0981(85)90240-0)
- 1056 Lapointe, B. E., & Ryther, J. H. (1978). Some aspects of the growth and yield of *Gracilaria*
 1057 *tikvahiae* in culture, *Aquaculture*, 15(3), 185-193. [https://doi.org/10.1016/0044-](https://doi.org/10.1016/0044-8486(78)90030-3)
 1058 [8486\(78\)90030-3](https://doi.org/10.1016/0044-8486(78)90030-3)
- 1059 Lee, M.-S., Wycislo, A., Guo, J., Lee, D. K., Voight, T. (2017). Nitrogen Fertilization Effects on
 1060 Biomass Production and Yield Components of *Miscanthus × giganteus*. *Frontiers in Plant*
 1061 *Science* 18, 544. Retrieved from <https://doi.org/10.3389/fpls.2017.00544>
- 1062 Lombardelli, G., Pirone, R., Ruggeri, B. (2017). LCA analysis of different MSW treatment
 1063 approaches in the light of energy and sustainability perspectives, *Chemical Engineering*
 1064 *Transactions*, 57, 469-474. <https://doi.org/10.3303/CET1757079>
- 1065 Lucas, K., Capron, M. E., Blaylock, R., Chambers, M. D., Stewart, J. R., DiMarco, S. F., et al.
 1066 (2019a). Ocean Forests: Feeding the World with Floating Artificial Reefs. *World*
 1067 *Aquaculture Society*. Retrieved from
 1068 <https://www.was.org/meetings/ShowAbstract.aspx?Id=136247>
- 1069 Lucas, K., Capron, M. E., Blaylock, R., Stewart, J. R., Howden, S., Chambers M. D., et al.
 1070 (2019b). AdjustaDepth Phase 1 Final Report to ARPA-E, DE-AR0000916.
 1071 <https://doi.org/10.13140/RG.2.2.29150.46405>
- 1072 Markets Insider. (2020). Oil Price of WTI Crude (West Texas Intermediate) History from 2006
 1073 to 2020. Retrieved from <https://markets.businessinsider.com/commodities/oil-price?type=wti>

- 1074 Martin, A. R., Erickson, D. L., Kress, W. J., & Thomas, S. C. (2014). Wood nitrogen
 1075 concentrations in tropical trees: phylogenetic patterns and ecological correlates. *New*
 1076 *Phytologist* 204: 484–495. <https://doi.org/10.1111/nph.12943>
- 1077 Matter, J. M., Stute, M., Snæbjörnsdóttir, S. Ó., Oelkers, E. H., Gislason, S. R., Aradóttir, E. S.,
 1078 et al. (2016). Rapid carbon mineralization for permanent disposal of anthropogenic carbon
 1079 dioxide emissions. *Science* 352, 1312–1314. <https://doi.org/10.1126/science.aad8132>
- 1080 McGlashan, N., Shah, N., Caldecott, B., & Workman, M. (2012). High-level techno-economic
 1081 assessment of negative emissions technologies. *Process Safety and Environmental*
 1082 *Protection*, 90(6), 501–510. <https://doi.org/10.1016/J.PSEP.2012.10.004>
- 1083 McGrail, B. P., Schaef, H. T., Spane, F. A., Horner, J. A., Owen, A. T., Cliff, J. B., Qafoku, O.,
 1084 Thompson, C. J., & Sullivan, E. C. (2017). Wallula Basalt Pilot Demonstration Project: Post-
 1085 injection Results and Conclusions. *Energy Procedia*, 114, 5783–5790.
 1086 <https://doi.org/10.1016/J.EGYPRO.2017.03.1716>
- 1087 McLaren, D. (2012). A comparative global assessment of potential negative emissions
 1088 technologies. *Process Safety and Environmental Protection*, 90(6), 489–500.
 1089 <https://doi.org/10.1016/J.PSEP.2012.10.005>
- 1090 McMahan, J. (2019). Multiple Zero-Carbon Natural-Gas Plants Planned At Lower Cost Than
 1091 Conventional Plants. *Forbes Magazine*. Retrieved from
 1092 [https://www.forbes.com/sites/jeffmcmahan/2019/07/23/net-power-developing-multiple-zero-](https://www.forbes.com/sites/jeffmcmahan/2019/07/23/net-power-developing-multiple-zero-carbon-natural-gas-plants-exec-says/#5762b0a68ac2)
 1093 [carbon-natural-gas-plants-exec-says/#5762b0a68ac2](https://www.forbes.com/sites/jeffmcmahan/2019/07/23/net-power-developing-multiple-zero-carbon-natural-gas-plants-exec-says/#5762b0a68ac2)
- 1094 Melzer, S. (2012). *Carbon Dioxide Enhanced Oil Recovery (CO₂ EOR): Factors Involved in*
 1095 *Adding Carbon Capture, Utilization and Storage (CCUS) to Enhanced Oil Recovery*. Report
 1096 for the National Enhanced Oil Recovery Initiative, Center for Climate and Energy Solutions.
 1097 Retrieved from [https://carboncapturecoalition.org/wp-](https://carboncapturecoalition.org/wp-content/uploads/2018/01/Melzer_CO2EOR_CCUS_Feb2012.pdf)
 1098 [content/uploads/2018/01/Melzer_CO2EOR_CCUS_Feb2012.pdf](https://carboncapturecoalition.org/wp-content/uploads/2018/01/Melzer_CO2EOR_CCUS_Feb2012.pdf)
- 1099 Metcalfe, G. E. (2020). What Should Carbon Cost?" *Scientific American* 322, 6, 62-69 (June
 1100 2020). doi:10.1038/scientificamerican0620-62
- 1101 Milledge, J.J. & Harvey, P.J. (2016), Potential process ‘hurdles’ in the use of macroalgae as
 1102 feedstock for biofuel production in the British Isles. *Journal of Chemical Technology and*
 1103 *Biotechnology* 91: 2221-2234. <https://doi.org/10.1002/jctb.5003>
- 1104 Minx, J. C., Lamb, W. F., Callaghan, M. W., Fuss, S., Hilaire, J., Creutzig, F., et al. (2018).
 1105 Negative emissions—Part 1: Research landscape and synthesis. *Environmental Research*
 1106 *Letters*, 13(6), 63001. <https://doi.org/10.1088/1748-9326/aabf9b>
- 1107 Monterey Bay Aquarium Research Institute (MBARI). (2019). Ocean Data FloatViz 6.0.
 1108 Retrieved from [https://www.mbari.org/science/upper-ocean-systems/chemical-sensor-](https://www.mbari.org/science/upper-ocean-systems/chemical-sensor-group/floatviz)
 1109 [group/floatviz](https://www.mbari.org/science/upper-ocean-systems/chemical-sensor-group/floatviz)
- 1110 Moran, K, Crawford, C, Webb, R, Holmes, G, Paulson, M, Rankin, M, et al. (2020). *Solid*
 1111 *Carbon: A Climate Mitigation Partnership Advancing Stable Negative Emissions*. Retrieved
 1112 from <https://www.solidcarbon.ca/>
- 1113 Morley, J. W., Selden, R. L., Latour, R. J., Frölicher, T. L., Seagraves, R. J., & Pinsky, M. L.
 1114 (2018). Projecting shifts in thermal habitat for 686 species on the North American continental
 1115 shelf. *PLOS ONE*, 13(5), e0196127. <https://doi.org/10.1371/journal.pone.0196127>

- 1116 Müller, R. D., Seton, M., Zahirovic, S., Williams, S. E., Matthews, K. J., Wright, N. M., et al.
 1117 (2016). Ocean basin evolution and global-scale plate reorganization events since Pangea
 1118 breakup. *Annual Reviews of Earth and Planetary Science* **44**, 107–138.
- 1119 N'Yeurt, A. de R., Chynoweth, D. P., Capron, M. E., Stewart, J. R., & Hasan, M. A. (2012).
 1120 Negative carbon via Ocean Afforestation. *Process Safety and Environmental Protection*,
 1121 *90*(6), 467–474. <https://doi.org/10.1016/J.PSEP.2012.10.008>
- 1122 Nagabhushan, D., & Thompson, J. (2019) Carbon Capture & Storage in the United States Power
 1123 Sector. Report for the Clean Air Task Force. Retrieved from
 1124 <https://www.catf.us/resource/45q-ccs-analysis>
- 1125 National Academies of Sciences, Engineering, & Medicine. (2019). *Negative Emissions*
 1126 *Technologies and Reliable Sequestration: A Research Agenda*. The National Academies
 1127 Press. <https://doi.org/10.17226/25259>
- 1128 Nemet, G. F., Callaghan, M. W., Creutzig, F., Fuss, S., Hartmann, J., Hilaire, J., Lamb, W. F.,
 1129 Minx, J. C., Rogers, S., & Smith, P. (2018). Negative emissions—Part 3: Innovation and
 1130 upscaling. *Environmental Research Letters*, *13*(6), 63003. [https://doi.org/10.1088/1748-](https://doi.org/10.1088/1748-9326/aabff4)
 1131 [9326/aabff4](https://doi.org/10.1088/1748-9326/aabff4)
- 1132 Oelkers, E. H., Gislason, S. R., & Matter, J. (2008). Mineral Carbonation of CO₂. *Elements*, *4*(5),
 1133 333–337. <https://doi.org/10.2113/gselements.4.5.333>
- 1134 Otto, I. M., Donges, J. F., Cremades, R., Bhowmik, A., Hewitt, R. J., Lucht, W., et al. (2020).
 1135 Social tipping dynamics for stabilizing Earth's climate by 2050. *Proceedings of the National*
 1136 *Academy of Sciences*, *117*(5), 2354 LP – 2365. <https://doi.org/10.1073/pnas.1900577117>
- 1137 Pachauri, R. K. & Meyer, L. (2014). *Climate Change 2014: Synthesis Report. Contribution of*
 1138 *Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel*
 1139 *on Climate Change* (IPCC, Geneva, Switzerland). Retrieved from
 1140 <https://www.ipcc.ch/report/ar5/syr/>
- 1141 Pan, W. L., Port, L. E., Xiao, Y., Bary, A. I., & Cogger, C. G. (2017). Soil Carbon and Nitrogen
 1142 Fraction Accumulation with Long-Term Biosolids Applications. *Soil Science Society of*
 1143 *America Journal*, *81*, 1381–1388. <https://doi.org/10.2136/sssaj2017.03.0075>
- 1144 Pandur, Z., Šušnjar, M., Zorić, M., Nevečerel, H., and Horvat, D. (2015). Energy Return on
 1145 Investment (EROI) of Different Wood Products. In M. Zlatic (Ed.), *Precious Forests -*
 1146 *Precious Earth*. IntertechOpen. <https://doi.org/10.5772/61144>
- 1147 Paraskova, T. (2019). Global LNG Markets Are Circling The Drain. Retrieved from
 1148 <https://oilprice.com/Energy/Natural-Gas/Global-LNG-Markets-Are-Circling-The-Drain.html>
- 1149 Park, M., Shin, S. K., Do, Y. H., Yarish, C., & Kim, J. K. (2018). Application of open water
 1150 integrated multi-trophic aquaculture to intensive monoculture: A review of the current status
 1151 and challenges in Korea. *Aquaculture*, *497*, 174–183.
 1152 <https://doi.org/10.1016/J.AQUACULTURE.2018.07.051>
- 1153 Pauly, D. (2019). Micronutrient richness of global fish catches. *Nature*, *574*(7776), 41–42.
 1154 <https://doi.org/d41586-019-02810-2>
- 1155 Penniman, C. A. (1983). Ecology of *Gracilaria tikvahiae* McLachlan (Gigartinales, Rhodophyta)
 1156 in the Great Bay Estuary, New Hampshire. University of New Hampshire Doctoral
 1157 Dissertation. Retrieved from <https://scholars.unh.edu/dissertation/1405>

- 1158 Pichach, C. (2019). CleanCarbon Energy: Carbon Balanced Renewable Energy, plus private
1159 communication. Retrieved from <https://www.f6s.com/cleancarbonenergy>
- 1160 Poore, J., & Nemecek, T. (2018). Reducing food's environmental impacts through producers and
1161 consumers. *Science*, 360 (6392), 987–992. <https://doi.org/10.1126/science.aaq0216>
- 1162 Pope Francis. (2015). *Encyclical Letter Laudato Si' of the Holy Father Francis on Care for Our*
1163 *Common Home*. Retrieved from
1164 [http://www.vatican.va/content/francesco/en/encyclicals/documents/papa-](http://www.vatican.va/content/francesco/en/encyclicals/documents/papa-francesco_20150524_enciclica-laudato-si.html)
1165 [francesco_20150524_enciclica-laudato-si.html](http://www.vatican.va/content/francesco/en/encyclicals/documents/papa-francesco_20150524_enciclica-laudato-si.html)
- 1166 PRNewswire. (2018). NET Power Achieves Major Milestone for Carbon Capture with
1167 Demonstration Plant First Fire. Retrieved from [https://www.prnewswire.com/news-](https://www.prnewswire.com/news-releases/net-power-achieves-major-milestone-for-carbon-capture-with-demonstration-plant-first-fire-300656175.html)
1168 [releases/net-power-achieves-major-milestone-for-carbon-capture-with-demonstration-plant-](https://www.prnewswire.com/news-releases/net-power-achieves-major-milestone-for-carbon-capture-with-demonstration-plant-first-fire-300656175.html)
1169 [first-fire-300656175.html](https://www.prnewswire.com/news-releases/net-power-achieves-major-milestone-for-carbon-capture-with-demonstration-plant-first-fire-300656175.html)
- 1170 Quirion, P., Rozenberg, J., Sassi, O., Vogt-Schilb, A. (2011). How CO₂ Capture and Storage Can
1171 Mitigate Carbon Leakage. Fondazione Eni Enrico Mattei Working Paper 2011.015.
1172 Retrieved from [https://www.feem.it/en/publications/feem-working-papers-note-di-lavoro-](https://www.feem.it/en/publications/feem-working-papers-note-di-lavoro-series/how-co2-capture-and-storage-can-mitigate-carbon-leakage/)
1173 [series/how-co2-capture-and-storage-can-mitigate-carbon-leakage/](https://www.feem.it/en/publications/feem-working-papers-note-di-lavoro-series/how-co2-capture-and-storage-can-mitigate-carbon-leakage/)
- 1174 Radulovich, R., Umanzor, S., Cabrera, R., & Mata, R. (2015). Tropical seaweeds for human
1175 food, their cultivation and its effect on biodiversity enrichment. *Aquaculture*, 436, 40–46.
1176 <https://doi.org/10.1016/J.AQUACULTURE.2014.10.032>
- 1177 Ravindra, K., Singh, T., & Mor, S. (2019). Emissions of air pollutants from primary crop residue
1178 burning in India and their mitigation strategies for cleaner emissions. *Journal of Cleaner*
1179 *Production*, 208, 261–273. <https://doi.org/https://doi.org/10.1016/j.jclepro.2018.10.031>
- 1180 Reinhart, D. (2004). Estimation of Energy Content of Municipal Solid Waste. Retrieved from
1181 <http://msw.cecs.ucf.edu/wastecharacterization.html>
- 1182 REN21. (2019). *Renewables 2019 Global Status Report*. Retrieved from
1183 <https://www.ren21.net/gsr-2019/>
- 1184 ReNew ELP. (2019). *Unlocking the Value in Plastic Waste*. Retrieved from
1185 <https://renewelp.co.uk/>
- 1186 Ries, J. B. (2010). Review: geological and experimental evidence for secular variation in
1187 seawater Mg/Ca (calcite-aragonite seas) and its effects on marine biological calcification.
1188 *Biogeosciences*, 7(9), 2795–2849. <https://doi.org/10.5194/bg-7-2795-2010>
- 1189 Rivers, P. (2017). The density of corrugated cardboard. Retrieved from
1190 <https://www.hunker.com/13419984/the-density-of-corrugated-cardboard>
- 1191 Rockström, J., Gaffney, O., Rogelj, J., Meinshausen, M., Nakicenovic, N., & Schellnhuber, H. J.
1192 (2017). A roadmap for rapid decarbonization. *Science*, 355(6331), 1269–1271.
1193 <https://doi.org/10.1126/science.aah3443>
- 1194 Rogelj, J., Shindell, D., Jiang, K., Fifita, S., Forster, P., Ginzburg, V., et al. (2018). Mitigation
1195 pathways compatible with 1.5°C in the context of sustainable development. In Masson-
1196 Delmotte, V., Zhai, P., Pörtner, H.-O., Roberts, D., Skea, J., Shukla, P. R., et al. (eds.),
1197 *Global Warming of 1.5°C: An IPCC Special Report on the impacts of global warming of*
1198 *1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in*
1199 *the context of strengthening the global response to the threat of climate change, sustainable*

- 1200 *development, and efforts to eradicate poverty.* (pp. 1–82). Retrieved from
 1201 https://report.ipcc.ch/sr15/pdf/sr15_chapter2.pdf
- 1202 Rose, C., Parker, A., Jefferson, B., & Cartmell, E. (2015). The Characterization of Feces and
 1203 Urine: A Review of the Literature to Inform Advanced Treatment Technology. *Critical*
 1204 *Reviews of Environmental Science and Technology*, 2;45(17):1827-1879.
 1205 <https://doi.org/10.1080/10643389.2014.1000761>
- 1206 Rubin, E. S., Davison, J. E., & Herzog, H. J. (2015). The cost of CO₂ capture and storage.
 1207 *International Journal of Greenhouse Gas Control*, 40, 378–400.
 1208 <https://doi.org/10.1016/J.IJGGC.2015.05.018>
- 1209 Saini, J. K., Saini, R., & Tewari, L. (2015). Lignocellulosic agriculture wastes as biomass
 1210 feedstocks for second-generation bioethanol production: concepts and recent developments.
 1211 *3 Biotech*, 5(4), 337–353. <https://doi.org/10.1007/s13205-014-0246-5>
- 1212 Schulz, K. G., Hartley, S., & Eyre, B. (2019). Upwelling Amplifies Ocean Acidification on the
 1213 East Australian Shelf: Implications for Marine Ecosystems. In *Frontiers in Marine Science*,
 1214 6, p. 636). <https://doi.org/10.3389/fmars.2019.00636>
- 1215 Sepulveda, N. A., Jenkins, J. D., de Sisternes, F. J., Lester, R. K. (2018). Role of firm low-
 1216 carbon electricity resources in deep decarbonization of power generation. *Joule* 2, 2403–
 1217 2420. <https://doi.org/10.1016/j.joule.2018.08.006>
- 1218 Shahbandeh, M. (2019). Global Meat Industry - Statistics & Facts. Retrieved from
 1219 <https://www.statista.com/topics/4880/global-meat-industry/>
- 1220 Sherman, M. T., Lucas, K., & Stewart, J. R. (2018). SeaweedPaddock : Initial Modeling and
 1221 Design for a Sargassum Ranch. *OCEANS 2018 MTS/IEEE Charleston*, 1–6.
 1222 <https://doi.org/10.1109/OCEANS.2018.8604848>
- 1223 Shi, H., Zheng, W., Zhang, X., Zhu, M., & Ding, D. (2013). Ecological–economic assessment of
 1224 monoculture and integrated multi-trophic aquaculture in Sanggou Bay of China. *Aquaculture*,
 1225 410–411, 172–178. <https://doi.org/10.1016/J.AQUACULTURE.2013.06.033>
- 1226 Shrestha, P. (2020). UK first: Hydrogen injected into gas grid as landmark trial gets underway.
 1227 Retrieved from <https://www.energylivenews.com/2020/01/06/uk-first-hydrogen-injected-into-gas-grid-as-landmark-trial-gets-underway/>
 1228
- 1229 Sims, N. A., Vollbrecht, L., Bullock, M., & Martin, K. (2019). *Ocean Era Blue Fields Offshore*
 1230 *Macroalgae Demonstration Project*. Retrieved from <http://ocean-era.com/projects>
- 1231 Smeets, E. M. W., Faaij, A. P. C., Lewandowski, I. M., & Turkenburg, W. C. (2007). A bottom-
 1232 up assessment and review of global bio-energy potentials to 2050. *Progress in Energy and*
 1233 *Combustion Science*, 33(1), 56–106. <https://doi.org/10.1016/j.pecs.2006.08.001>
- 1234 Smil, V. (1999). Crop Residues: Agriculture’s Largest Harvest: Crop residues incorporate more
 1235 than half of the world’s agricultural phytomass. *BioScience*, 49(4), 299–308.
 1236 <https://doi.org/10.2307/1313613>
- 1237 Snæbjörnsdóttir, S. Ó., Sigfússon, B., Marieni, C., Goldberg, D., Gislason, S. R., & Oelkers, E.
 1238 H. (2020). Carbon dioxide storage through mineral carbonation. *Nature Reviews Earth &*
 1239 *Environment*, 1(2), 90–102. <https://doi.org/10.1038/s43017-019-0011-8>

- 1240 Sorondo, M. S., & Ramanathan, V. (2016). Pursuit of integral ecology. *Science*, 352(6287), 747.
 1241 <https://doi.org/10.1126/science.aag0826>
- 1242 StatisticsTimes. (2019). *Gross World Product*. Retrieved from
 1243 <http://statisticstimes.com/economy/gross-world-product.php>
- 1244 Steeper Energy. (2019). *Hydrofaction technology*. Retrieved from <https://steeperenergy.com/>
- 1245 Strauss, W. & Walker, S. (2018). Global pellet market outlook in 2018, Canadian Biomass
 1246 Magazine, 2 February 2018. Retrieved from
 1247 <https://www.canadianbiomassmagazine.ca/pellets/increasing-demand-6705>
- 1248 Sumaila, U. R., Tai, T. C., Lam, V. W. Y., Cheung, W. W. L., Bailey, M., Cisneros-
 1249 Montemayor, A. M., et al. (2019). Benefits of the Paris Agreement to ocean life, economies,
 1250 and people. *Science Advances*, 5(2), eaau3855. <https://doi.org/10.1126/sciadv.aau3855>
- 1251 Sverdrup, H. U., Johnson, M. W., Fleming, R. H. (1942). *The Oceans, Their Physics, Chemistry,*
 1252 *and General Biology*. New York: Prentice-Hall. Retrieved from
 1253 <http://ark.cdlib.org/ark:/13030/kt167nb66r/>
- 1254 Tàbara, D., Frantzeskaki, N., Hölscher, K., Pedde, S., Kok, K., Lamperti, F., Christensen, J.,
 1255 Jäger, J., & Berry, P. (2018). Positive tipping points in a rapidly warming world. *Current*
 1256 *Opinion in Environmental Sustainability*, 31, 120–129.
 1257 <https://doi.org/10.1016/j.cosust.2018.01.012>
- 1258 Tang, Z., Xu, W., Zhou, G., Shu, J., Han, W., Bai, Y., et al. (2014). Patterns of plant carbon,
 1259 nitrogen, and phosphorus concentration in relation to productivity in China's terrestrial
 1260 ecosystems. *PNAS* April 17, 2018 115 (16) 4033-4038.
 1261 <https://doi.org/10.1073/pnas.1700295114>
- 1262 Tanager P, Field J L, Jahn C E, DeFoort M W, Leach J E. (2013) Biomass for thermochemical
 1263 conversion: targets and challenges. *Frontiers in Plant Science* 4, Article 218.
 1264 doi:10.3389/fpls.2013.00218
- 1265 TechLink. (2019). The 9 Technology Readiness Levels of the DoD. Retrieved from
 1266 <https://techlinkcenter.org/technology-readiness-level-dod>
- 1267 Timko, M. T., Ghoniem, A. F., Green, W. H. (2015). Upgrading and desulfurization of heavy
 1268 oils by supercritical water. *The Journal of Supercritical Fluids*, 96, 114–123.
 1269 <https://doi.org/10.1016/j.supflu.2014.09.015>
- 1270 The Royal Society. 2020. Ammonia: zero-carbon fertiliser, fuel and energy store. February 2020
 1271 DES5711 ISBN: 978-1-78252-448-9. Retrieved from [https://royalsociety.org/topics-](https://royalsociety.org/topics-policy/projects/low-carbon-energy-programme/green-ammonia/)
 1272 [policy/projects/low-carbon-energy-programme/green-ammonia/](https://royalsociety.org/topics-policy/projects/low-carbon-energy-programme/green-ammonia/)
- 1273 Turner, P. A., Mach, K. J., Lobell, D. B., Benson, S. M., Baik, E., Sanchez, D. L., et al. (2018).
 1274 The global overlap of bioenergy and carbon sequestration potential. *Climatic Change*,
 1275 148(1), 1–10. <https://doi.org/10.1007/s10584-018-2189-z>
- 1276 Tvinnereim, E., & Mehling, M. (2018). Carbon pricing and deep decarbonisation. *Energy Policy*,
 1277 121, 185–189. <https://doi.org/10.1016/J.ENPOL.2018.06.020>
- 1278 U.S. Department of Energy Advanced Research Projects Agency-Energy. (2017a). MacroAlgae
 1279 Research Inspiring Novel Energy Resources (MARINER) Project Listing. Retrieved from
 1280 <https://arpa-e.energy.gov/?q=arpa-e-programs/mariner>

- 1281 U.S. Department of Energy Advanced Research Projects Agency-Energy. (2017b). *MacroAlgae*
 1282 *Research Inspiring Novel Energy Resources (MARINER) Program Overview*. Retrieved from
 1283 <https://arpa->
 1284 [e.energy.gov/sites/default/files/documents/files/MARINER_ProgramOverview_FINAL.pdf](https://arpa-e.energy.gov/sites/default/files/documents/files/MARINER_ProgramOverview_FINAL.pdf)
- 1285 U.S. Department of Energy. (2018). *2016 Billion-Ton Report: Advancing Domestic Resources*
 1286 *for a Thriving Bioeconomy*. <https://doi.org/10.2172/1271651>
- 1287 U.S. Department of Energy. (2019). *U.S. Department of Energy Invests \$7 Million for Projects*
 1288 *to Advance Coal Power Generation under Coal FIRST Initiative*.
 1289 [https://www.energy.gov/fe/articles/us-department-energy-invests-7-million-projects-](https://www.energy.gov/fe/articles/us-department-energy-invests-7-million-projects-advance-coal-power-generation-under-coal)
 1290 [advance-coal-power-generation-under-coal](https://www.energy.gov/fe/articles/us-department-energy-invests-7-million-projects-advance-coal-power-generation-under-coal)
- 1291 U.S. Department of Energy. (2020). Reports for the Coal FIRST concepts of interest. National
 1292 Energy Technology Laboratory. Retrieved from
 1293 <https://netl.doe.gov/coal/tpg/coalfirst/concept-reports>
- 1294 U.S. Energy Information Administration (EIA). (2005). Average Heat Content of Selected
 1295 Biomass Fuels. Renewable Energy Annual 2004. Retrieved from
 1296 <https://www.eia.gov/renewable/annual/>
- 1297 U.S. Energy Information Administration. (2019). Coal explained: Coal prices and outlook.
 1298 <https://www.eia.gov/energyexplained/coal/prices-and-outlook.php>
- 1299 U.S. Energy Information Administration. (2020). Petroleum and other liquids.
 1300 [https://www.eia.gov/international/data/world/petroleum-and-other-liquids/annual-petroleum-](https://www.eia.gov/international/data/world/petroleum-and-other-liquids/annual-petroleum-and-other-liquids-production)
 1301 [and-other-liquids-production](https://www.eia.gov/international/data/world/petroleum-and-other-liquids/annual-petroleum-and-other-liquids-production)
- 1302 UNEP. (2019a). Emissions Gap Report 2019. Retrieved from
 1303 <https://www.unenvironment.org/resources/emissions-gap-report-2019>
- 1304 UNEP. (2019b). Waste-to-Energy: Considerations for Informed Decision-Making. Retrieved
 1305 from <http://wedocs.unep.org/bitstream/handle/20.500.11822/28413/WTEfull.pdf>
- 1306 UNFCCC. 2015. What is the Paris Agreement? Retrieved from [https://unfccc.int/process-and-](https://unfccc.int/process-and-meetings/the-paris-agreement/what-is-the-paris-agreement)
 1307 [meetings/the-paris-agreement/what-is-the-paris-agreement](https://unfccc.int/process-and-meetings/the-paris-agreement/what-is-the-paris-agreement) (accessed April 14, 2020).
- 1308 United Nations. (2016a). Agenda for Humanity. Retrieved from
 1309 <https://www.agendaforhumanity.org>
- 1310 United Nations. (2016b). Sustainable Development Goals. Retrieved from
 1311 <https://sustainabledevelopment.un.org>
- 1312 USGCRP. (2018). Impacts, Risks, and Adaptation in the United States: Fourth National Climate
 1313 Assessment, Volume II. (B. C. S. Reidmiller, D.R., C.W. Avery, D.R. Easterling, K.E.
 1314 Kunkel, K.L.M. Lewis, T.K. Maycock (ed.)). <https://doi.org/10.7930/NCA4.2018>.
- 1315 Vance E. (2017). Baja California's recipe for saving fishing communities. *National Geographic*
 1316 September 2017. Retrieved from
 1317 [https://www.nationalgeographic.com/magazine/2017/09/baja-mexico-marine-conservation-](https://www.nationalgeographic.com/magazine/2017/09/baja-mexico-marine-conservation-tourism-fish-sharks-whales/)
 1318 [tourism-fish-sharks-whales/](https://www.nationalgeographic.com/magazine/2017/09/baja-mexico-marine-conservation-tourism-fish-sharks-whales/)
- 1319 Wallace, M., Kuuskraa, V., & DiPietro, P. (2013). CO2 Utilization from "Next Generation" CO2
 1320 Enhanced Oil Recovery Technology. *Energy Procedia* 37:6854-6866.
 1321 <https://doi.org/10.1016/j.egypro.2013.06.618>

- 1322 Watson, J., Wang, T., Si, B., Chen, W. T., Aierzhati, A., & Zhang, Y. (2020). Valorization of
 1323 hydrothermal liquefaction aqueous phase: pathways towards commercial viability. *Progress*
 1324 *in Energy and Combustion Science*, 77. <https://doi.org/10.1016/j.pecs.2019.100819>
- 1325 Wennerholm, M. (2020). AquaDam Special Project Applications. Retrieved from
 1326 <https://www.aquadam.net>
- 1327 Wessler, S. (2019). *In King County, Washington, human waste is a climate solution*. Retrieved
 1328 from [https://www.yaleclimateconnections.org/2019/11/in-king-county-washington-human-](https://www.yaleclimateconnections.org/2019/11/in-king-county-washington-human-waste-is-a-climate-solution/)
 1329 [waste-is-a-climate-solution/](https://www.yaleclimateconnections.org/2019/11/in-king-county-washington-human-waste-is-a-climate-solution/)
- 1330 West, J. J., Smith, S. J., Silva, R. A., Naik, V., Zhang, Y., Adelman, Z., et al. (2013). Co-benefits
 1331 of mitigating global greenhouse gas emissions for future air quality and human health.
 1332 *Nature Climate Change*, 3(10), 885–889. <https://doi.org/10.1038/nclimate2009>
- 1333 Whittaker, JM, Alfonso JC, Masterton S, Müller D, Wessel P, Williams S, et al. (2015). Long-
 1334 term interaction between mid-ocean ridges and mantle plumes. *Nat. Geosci.* 8, 479–483.
 1335 <https://doi.org/10.1038/NGEO2437>
- 1336 Wilkinson K. 2020. The Drawdown Review 2020: Climate Solutions for a New Decade.
 1337 Retrieved from <https://drawdown.org/drawdown-framework/drawdown-review-2020>
 1338 (accessed April 10, 2020).
- 1339 Wilkinson, K. (2020). *The Drawdown Review 2020: Climate Solutions for a New Decade*.
 1340 <https://drawdown.org/drawdown-framework/drawdown-review-2020>
- 1341 Wood, D., Capuzzo, E., Kirby, D., Mooney-McAuley, K., & Kerrison, P. (2017). UK
 1342 macroalgae aquaculture: What are the key environmental and licensing considerations?
 1343 *Marine Policy*, 83(August 2016), 29–39. <https://doi.org/10.1016/j.marpol.2017.05.021>
- 1344 Yang, Y., Hagerly, M., Palmarozzo, A., Celebi, M., Chupka, M., & Sheffield, H. (2017). The
 1345 Future of Cap-and-Trade Program in California: Will Low GHG Prices Last Forever?
 1346 [https://brattlefiles.blob.core.windows.net/files/11768_the_future_of_cap-and-](https://brattlefiles.blob.core.windows.net/files/11768_the_future_of_cap-and-trade_program_in_california_final_12.4.17.pdf)
 1347 [trade_program_in_california_final_12.4.17.pdf](https://brattlefiles.blob.core.windows.net/files/11768_the_future_of_cap-and-trade_program_in_california_final_12.4.17.pdf)
- 1348 Zaines, G. G., & Khanna, V. (2013). Microalgal biomass production pathways: evaluation of
 1349 life cycle environmental impacts. *Biotechnology for Biofuels*, 6(1), 88.
 1350 <https://doi.org/10.1186/1754-6834-6-88>
- 1351