

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/303862554>

Environmental eco-friendly marine resource macro algae (Seaweeds): an omnipotent source for value added products and its applications – A review

Research · June 2016

DOI: 10.13140/RG.2.1.4024.5363

CITATIONS

0

READS

803

1 author:



Dr. B. Bharathiraja

Veltech High Tech Dr.Rangarajan Dr.Sakunthala Engineering College

148 PUBLICATIONS 1,330 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Microbial production of high value Malic acid from low cost crude glycerol [View project](#)



Microbial production of high value malic acid from low cost crude glycerol [View project](#)

Review Article

doi: <http://dx.doi.org/10.20546/ijcmas>

Environmental eco-friendly marine resource macro algae (Seaweeds): an omnipotent source for value added products and its applications - A review

B.Bharathiraja^{1*}, P.Devaki¹, S.Dheepa¹, R.Mageshwari¹, J. Jayamuthunagai²,
M. Chakravarthy¹, D. Yuvaraj¹ and R. Praveenkumar^{3*}

¹Vel Tech High Tech Dr. Rangarajan Dr. Sakunthala Engineering College, Avadi,
Chennai – 600062, Tamilnadu, India

²Centre For Biotechnology, Anna University, Chennai, Tamilnadu-600025, India

³Department of Biotechnology, Arunai Engineering College,
Tiruvannamalai–606603, Tamilnadu, India

*Corresponding author email id: btrbio@gmail.com

ABSTRACT

Keywords

Seaweed,
Nutritious food,
Biofuels, Bioactive
compounds,
Polysaccharides,
Nanoparticles.

Article Info

Accepted:
xx June 2016
Available Online:
xx July 2016

The consequent reduction of land resources by human activities such as pollution, over exploitation, industrialization, migration etc., has lead man to search for other alternative ways to meet the demands for well-being. When this has been the existing situation in most of the developing countries, kelp forest resources like seaweeds which are known for its potentially strong bioactive compounds can be the best fit to fulfill numerous requirements such as nutritious food, biofuels, biofertilizer and pharmaceuticals to cure different diseases and other industrial applications. From the history till date seaweed has been employed in various sectors like food, pharmaceuticals, agriculture, waste water treatment and so on. Malnutrition and poverty have become the major issue in many of the developing nations, hale and healthy life can be guaranteed by seaweed food products and also people can be self- employed. Besides many research works done regarding seaweeds this review provides a collective idea about the potentials and wide range applications of seaweed, a marine macroalgae.

Introduction

Seaweed which is a macroalgae seen in various form, color and occur along the coastal line in marine habitat and India as a peninsular nation has got wide distribution of seaweeds compared to other countries (Brian E. Lapointe *et al.*, 1981, Subba Rao, *et al.*, 2006). Several harvesting techniques and cultivation methods for red, brown and green algae were in practice depending upon the species and other environmental factors (Buschmann *et al.*, 1995; Westermeier *et al.*,

1991). The biochemical components of seaweed such as carbohydrates, proteins, vitamins, fat, mineral are numerous and their composition experiences seasonal variation. The various compounds such as sulfated polysaccharides, fucoxanthin, carrageenans, agarans, terpenoids etc., and organic extracts such as methanolic, ethanolic and butanolic extracts of different seaweeds possess antioxidant, antiviral, anti-inflammatory, anticancer activity against

various pathogens and diseases. With the increased domestic, industrial and vehicle uses seaweed compounds extracted and processed are being used as various biofuels. Moreover *Ulva* species has been used in the production of all the four biofuels like bioethanol, biodiesel, biobutanol, biogas while biomass of other macroalgal species such as *Sargassum*, *Padina*, *Dictyota*, *Ascophyllum*, *Laminaria*, *Saccharina* were also found to be used in the production of various fuels. On the other hand polysaccharides of seaweeds are used to produce agar a phycocolloid which has its application in the food industries in the form of stabilizer, thickening agent; other carbohydrates such as fucoidan, laminarin, carrageenan were also analysed to have varied applications in the food and drugs sector. Seaweed extracts from *Laminaria*, *Undaria*, *Saccharina* species were known to be employed in cancer treatment whereas other mineral rich extracts treat arthritis and hypothyroidism. Current world which approaches the trends of nanotechnology, silver and gold nanoparticles from potential seaweed species was found to have intense antimicrobial properties that aids effluent treatment. Apart from these seaweeds such as *Laminaria*, *Sargassum*, *Ulva* species were being used in treating waste water, *Ascophyllum* species as beauty enhancers, nutritious animal feed while their alkaline extracts as biofertiliser have provided good quality plants.

Morphology and Diversity

Algae have a great variety of forms, sizes and colors in which some are simple colonies with many cells; some are filamentous, tubular, meshed, membranous or saccate algae. Some are more delicate and complex, such as, *Sargassum* having structures similar to roots, stems and leaves of plants as well as differentiated air bladders which help the algae floating to

water surface to be able to absorb more sunlight. Nevertheless, although algae may have different forms, their internal structures are composed of similar cells with simple differentiation instead of true roots, stems or leaves. Seaweeds are grouped into three based on colour: Green algae (Chlorophyta), brown algae (Phaeophyta), Red algae (Rhodophyta), Blue-green algae (Cyanophyta) which are being harvested and utilized for several purposes (Brian E. Lapointe *et al.*, 1981). The thallus (the algal body), lamina or blade (a flattened leaf-like structure), stipe (a stem-like structure), holdfast (a basal structure provides attachment to a surface) forms the parts of a seaweed which is shown in the figure 1.

Occurrence and Distribution

Seaweed is particularly available in abundant in three areas: the warm northern waters around the Kermadec and three king island, the Cook Strait Kaikoura coast region in central New Zealand and the south in an area encompassing Fiordland, Stewart island and the Otago coast. Also it was found that enumerable amount of varied species belonging to different genus are found along the Indian coast (Subba Rao, P. V *et al.*, 2006).

The figure 2 & 3 shows the different kinds of species distribution that includes Chlorophyta, Phaeophyta, Rhodophyta and Cyanophyta alongside countries of Indian ocean and areas of Indian coastal line (Vinod K. Dhargalkar., 2014) whereas figure 4 picturizes the seaweed strength along the coastal areas of other countries like China, Korea, Japan, USA, Ireland, Iceland, Scotland.

Cultivation and Harvesting

With the advent of various seaweed species cultivation of seaweed attracts more

attention to put seaweed in varied uses. The physical, chemical and biological parameters should be investigated before cultivation. Apart from this, the general environmental factors that influence the cultivation and harvesting processes are location, water and land quality, fauna, climatic factor, marketing accessibility, farm management operations, industrial and domestic pollution. Depending upon the properties and natural environment of individual seaweed species or species in groups, the harvesting methods are selected. The seaweed culture methods that are in practice are off-bottom method, raft method, horizontal culture method, mixed culture method, long line method. It is noted that the major problem faced by the seaweed industry is the lack of efficient harvesting technique. Let us see the viable harvesting that has been used from the past. The culturing method varies between species to species and for instance the red algae are generally seen in deep waters and are much smaller in size compared to the brown algae. In addition, the harvesting of red algae is expensive and involves complexity and it was reviewed that two planting techniques were commonly used for red algal farming. One is the direct method that involves the direct burial of the thalli into the sandy bottom using different types of tools and the other is the Plastic Tube Method which consists of fastening bundles of thalli to plastic tubes filled with sand, which anchor the algae to the sea bottom (Buschmann, A. H *et al.*, 1995). It was estimated from the experience gained from subtidal farms that hand-pulling of thalli has given more production rather than the use of tools (Westermeier *et al.*, 1991). Mechanized harvesting that uses power-driven barges equipped with reciprocating underwater mowers (cropping vessels) were used for cultivating *Macrocystis* that grows in large beds. Similar cropping equipments were

used to harvest *Ascophyllum* species which grows at high tide. The predominant species *Porphyra yezoensis*, *Porphyra tenera* grow in the inner parts of estuaries and bays and can survive in high salt conditions whereas *Porphyrapseudo linearis* grows in the deeper waters. The Concochelis–Spore culture technique was used to cultivate these *Porphyra*(nori) species (Suto, S., 1971). The rope cultivation and stone techniques were used in Japan to cultivate the brown alga *Undaria* or "wakame". The "collector strings" that are hung in the water for the sporophytes of *Undaria* to grow and tanks were used for seedling. As an expansion of past technique, along with natural beds a new substrate of rocks exploded with dynamite was also used to grow *Undaria* (Tamura, T., 1966). Production of hybrid varieties of *Undaria* was also made possible by crossing closely related species. To harvest *Laminaria* that grows only on hard, rocky ocean bottoms, reciprocating cutters mounted on dredges or a system of continuous grapnels were used and the development of techniques for cutting the normal 2-yr growth period to harvest down to 1yr was researched to increase the production (Hasegawa, Y., 1971). It was reported that in Norwegian coastal area the sugar kelp *Saccharina latissima* was cultivated by integrating with salmon (*Salmo salar*) aquaculture and the growth was seemed to be good in late autumn and in spring (Aleksander Handa Silje Forbord *et al.*, 2013).

Green algae may be found in marine or freshwater habitats, and some even thrive in moist soil. The green macro algae were cultivated in outdoor tanks and the biocrude was obtained through hydrothermal liquefaction in batch reactor. Also, the green algae were cultivated in ponds, open lagoons and in cages. It was reported that the *Oedogonium* species has given the

maximum yield followed by *Derbesia*, *Ulva* species (Neveux *et al.*, 2014).

Biochemical Composition

The composition of seaweeds varies depending upon season, density and other environmental factors. Compared to vegetables, fruits, pulses and cereals, seaweed records the maximum content of carbohydrates, proteins, vitamins, minerals, fat, fibre, ash, moisture. Earlier work has revealed that the different species gathered from similar area, family, environmental factors showed greater variations with respect to their elemental composition (Murugaiyan *et al.*, 2012). Also it has been observed that red seaweed contains thirty times more potassium than bananas, 200 times more iron than beetroot, the *Nori* seaweed constitute twice the protein than meat and the *Hijiki* seaweed contains twice the amount of calcium compared to full-cream milk. Further, it was noted that 15,000 novel compounds were chemically determined and algae are considered to be the predominant source for novel biologically active compounds that is required for human nutrition (Ivanova *et al.*, 2013). While seasonal variations are of great concern, it was showed that the phaeophyta and rhodophyta recorded the maximum biomass during summer and chlorophyta was maximum during autumn season (Dadollahi-Sohrab *et al.*, 2012). The table 1 shows the composition of some commercially important seaweeds.

Properties

It was reviewed that seaweeds are rich in bioactive compounds that promotes and some of the species to say are. *Laminaria* species, *Fucus* species, *Ascophyllum nodosum*, *Chondrus crispus*, *Porphyra* species, *Ulva* species, *Sargassum* species,

Gracilaria species and *Palmaria palmate* (Susan *et al.*, 2010). Since chemical preservatives were proved to cause deleterious health hazards like cancer, asthma etc., seaweeds have been known to be the safe and promising replace as food additives that has antimicrobial, antioxidant properties (Kasi Pandima Devi *et al.*, 2008). Also it was identified that many secondary metabolites from the marine source with effective antibacterial, antifungal and antiviral activities which are being used as antibiotics and drugs to treat various infectious diseases (Abad *et al.*, 2011). Bioactive compounds of marine macro algae are known to have diverse mechanism of action against diseases. In addition to this, red, brown and green macro algae were detected to contain compounds with cytostatic, antiviral, antihelminthic, antifungal and antibacterial activities (Newman *et al.*, 2003).

Antioxidant Activity

Algae are antioxidant in nature because of their non-enzymatic antioxidant components like reduced glutathione (GSH), ascorbic acid, phatocopherol, betacarotenoids, flavonoids, hydroines, pycocyanin, proline, mannitol, myoinositol, phenolics and polyamines (Fu-Jin Wang *et al.*, 2011). The antioxidants can inhibit the initiation of the oxidative chain reaction and thus prevent cell damage caused by Reactive Oxygen Species (ROS) (Sung-Myung Kang *et al.*, 2011). The antioxidant molecules can destroy free radicals by donating hydrogen atoms or by electron donation. In most of the cases DPPH had been used as a free radical to evaluate reducing substances rather than nitric oxide, deoxyribose, hydrogen peroxide, ABTS. The DPPH radical scavenging activity was evaluated using an ESR (Electron Spin Resonance) spectrometer (H. Indu and R. Seenivasan.,

2013). From the earlier works done it was reported that the extracts of species such as *Hijik fusiformis*, *Cladosiphon okamuranus*, *Undaria pinnatifida*, and *Sargassum fulvellum* were known to possess effective DPPH radical scavenging activity (Mise *et al.*, 2011, Yan *et al.*, 1999). Also Carotenoids have radical scavenging which helps to keep up health and in prevention of disease whereas Fucoxanthin has been reported to effectively scavenge chemically-generated free radicals like DPPH (W. Stahl and H. Sies., 2012). Based on quenching rate constants it was shown that the radical scavenging activity of fucoxanthin and its metabolite fucoxanthinol were higher than that of α -tocopherol and lower than β -carotene (K. Mikami and M. Hosokawa., 2013). Fucoxanthin (FX), an orange colored carotenoid belonging to non provitamin carotenoids which is also known as xanthophylls are present in edible brown seaweeds such as *Undaria pinnatifida*, *Hijikia fusiformis*, *Laminaria japonica* and *Sargassum fulvellum* possess significant antioxidant activity [99,100]. Sulfated polysaccharides (SP) are the compounds found in the extracellular matrix of seaweeds that has got antioxidant property and the well-known sp. are carragenans, agarans, xylase, galactose, mannose, fucan, fucoidan (Nednaldo *et al.*, 2012). Based on the rheological behavior, the polysaccharide content and the methanol extract of red algae *Gracillaria biridae* and *Gracilaria verrucosa* were reported to possess antioxidant property (De almeida *et al.*, 2011, Bartolomeu *et al.*, 2012). Further it was noted that the water extract of *Laminaria* species had higher antioxidant activity than its ethanol extract (Ismail A Jr and Tan S., 2002). Thus high levels of oxidative stress leads to many harmful diseases such as atherosclerosis, Parkinson's disease, Alzheimer's disease, acute myocardial infarction, chronic fatigue

syndrome and fibromyalgia. FX was estimated as an effective tool to prevent and treat these diseases (Nicolantonio *et al.*, 2012).

Anticancer Activity

Cancer being a fatal disease has become a major health problem worldwide mainly because of bad food habits. The conventional chemotherapy or radiotherapy treatment to control tumors and reduce the risk of mortality rates has resulted in other ill-effects such as long term side effects, destruction of healthy tissues etc. Several types of carcinomas such as prostate cancer, leukemia, colon cancer, breast cancer, liver cancer, melanoma and lymphoma are in existence (Thamaraiselvan *et al.*, 2013). In a quest to cure and uproot cancer, seaweeds were identified as the best source that has antitumor properties. The seaweed compounds or extracts undergo several types of mechanisms against the cancer cells and some of them are apoptosis induction, inhibition of tumor invasion, hyaluronidase activity inhibition, anti-angiogenic activity. Regulation of mammary gland integrity (Suhaila *et al.*, 2012). Polysaccharides present in the cell wall of macroalgae contain immunomodulatory and anticancer effects and are most considered in the medical areas of study (Shilpi *et al.*, 2011). Sulfated polysaccharides from brown species viz., *Sargassum*, *Laminaria*, *Ecklonia* inhibited growth of Sarcoma-180 cells and acted as antitumor against L-1210 leukemia (Yamamoto *et al.*, 1974, 1981, 1984a). Further fucoidan or fucose containing sulfated polysaccharides (FCSP) and MAPK (mitogen activated protein kinase) along with FCSP in brown seaweeds found to enhance and augment macrophage mediated immune signaling molecules production and thereby induced apoptosis (Marcel *et al.*, 2011). Breast cancer

ranks the second most common cancer in the world and is the major cause for mortality in women (Laura *et al.*, 2006). As a remedy for this, the methanol extract of *Sargassum muticum* activity against proliferation of breast cancer cell lines were evaluated for its apoptosis property (Farideh *et al.*, 2013). Also the combination of seaweed *Porphyra dentate*, β -sitosterol and campesterol was known to reduce the tumor size considerably (Ghislain *et al.*, 2014). Nextly the colorectal cancer which affects both men and women, their cancer cell lines proliferation could be inhibited by the extracts of Laminarian species and *Ulva fasciata* by apoptosis induction mechanism (Ryu *et al.*, 2013, Hee-Kyoung *et al.*, 2012, 2013). Furthermore based on dosage the anti-tumor and anti-metastatic activities of fucoidan isolated from *Fucus evanescens* were studied. Alginates and palmitic acid from the species *Sargassum vulgare* (brown algae) and *Amphiroa zonata* (red algae) were also reported to possess antitumor property (Alekseyenko *et al.*, 2007, Harada *et al.*, 2002).

Antimicrobial Activity

Algal biomass has been analysed to have antimicrobial activity and it mainly depends upon the algal species taken and the extraction method followed. The antimicrobial compound extracted from a biological source is generally by means of attacking the cell wall and cell membrane of the target organism. Further it disrupts the electron transport chain, coagulates protein and nucleic acid synthesis (Gupta, S. and N. Abu-Ghannam. 2011). Among seaweed species the ethyl acetate extracts of *Sargassum* species have strong antimicrobial activity against bacteria and fungi than the water extract and the activity is due to the presence of meroterpenoids (Horie *et al.*, 2008). The brown alga *Stoechospermum*

marginatum was active against bacterial strains *Klebsiella* and *Vibrio cholerae* whereas the green alga *Cladophora prolifera* was bacteriocidal against *Saccharomyces aureus* and *Vibrio cholera* (Muruleedhara *et al.*, 2003, Rodrigues *et al.*, 2004). It was found that the extracts of red algae *Gracilaria fisheri* inhibits the pathogen *Vibrio harveyi* which affects the Shrimp population (Kanjana *et al.*, 2011). Also the butanolic extracts of the seaweeds *Ulva lactuca* and *Sargassum wightii* exhibited considerable inhibition zone against the shrimp pathogen *Vibrio parahaemolyticus* (Immanuel *et al.*, 2004). The phlorotannins of *Ascophyllum nodosum* was more active against the *Escherichia coli* strain than the condensed or terrestrial tannins (Wang *et al.*, 2009). The dichloromethane extracts of several seaweed has showed significant antibacterial action against fish pathogens such as *Asparagopsis armata*, *Falkenbergia rufolanosa* (Bansemir *et al.*, 2006). The dimethyl sulfoxide (DMSO) extracts of seaweed species and their antiprotozoal activity against *Plasmodium* species has been studied and the selectivity index was the parameter used to evaluate the activity (Catherine *et al.*, 2011). The same DMSO extract of *Sargassum Longifolium* showed inhibitory activity against various bacterial strains whereas the acetic acid extract showed maximum inhibition against *Proteus* species and minimum activity against *Streptococcus* species (Ponnanikajamdeen *et al.*, 2014). The antifungal activity against *Phythium phanidermatum* and *Colletotrichum capsici* was maximum in *Ulva fasciata*. Ultimately nanoparticles which is the most welcomed compound because of its size and viability was found to be extracted as silver nanoparticles from *Ulva lactuca* by agar well diffusion method and it was analysed for its antibacterial activity (K. Sivakumar and T.Thinakaran., 2013).

Antiviral Activity

Promising feature of seaweed is its antiviral potential by which many fatal diseases can be treated. AIDS is a fatal disease caused by Human Immunodeficiency Virus (HIV) which belongs to retro virus family and has no effective treatment till date. The antiviral activity of seaweeds depends on the dosage and time. Apart from other causes the Herpes Simplex Virus (HSV-1) and HSV-2 which causes infection in mouth, face and genital area was also identified as a major risk factor for Human immunodeficiency Virus (HIV) (Vo *et al.*, 2011, Celum 2004) and the antiviral activity against these virus has been conferred with SQDS (Sulfoquinovosyldiacyl glycerol) fractions extracted from *Sargassum vulgare* (Erwan *et al.*, 2013). Also the bioactive alginates from *Sargassum* species has antiviral property and it was studied that the extracts of this species acted against the viruses Human T cell Lymphotropic Virus Type1 (HTLV1) and Human Immuno deficiency Virus Type1 (HIV-1) (Liu *et al.*, 2012, Romanos *et al.*, 2002, Mi-Jeong Ahn *et al.*, 2002). The fucoidan polysaccharide was against HIV and human cytomegalovirus (HCMV) while its derivative galactofuran extracted from the seaweed *Adenocystis utricularis* showed inhibitory action towards the retro viruses (HSV) 1 and 2 (Ponce *et al.*, 2003). Further the diterpenes isolated from the Dictyota species exerted antiviral action and galactofucan sulfate extract from *Undaria pinnatifida* worked against HSV-1, HSV-2 and HCMV (Human Cyto Megalovirus) (Siamopoulou *et al.*, 2004, Queiroz *et al.*, 2008). Considering plant viruses the seaweed polysaccharides such as fucans, laminarin, alginates, ulvans obtained from *Fucus vesiculosus*, *Laminaria digitatum*, *Lessonia* species, *Ulva* species respectively, when injected in to tobacco plants, protected against Tobacco Mosaic Virus (TMV) by

inducing Jasmonic acid (JA) and Salicylic acid (SA) signaling pathways (Jeannette *et al.*, 2011).

Anti-inflammatory activity

In general inflammatory response is an auto defensive mechanism that is met with huge leukocyte production and the inflammatory reactions are generally due to the presence of ROS, nitric oxide and other factors which results in tissue damage (Yong-Fang *et al.*, 2001). Caulerpin an alkaloid extracted from *Caulerpa racemosa* was found to exhibit anti-inflammatory properties (De Souza *et al.*, 2009) and this compound act by suppressing the antigen, histamine secretion, lymphocyte and natural killer cell proliferation (Mohamed *et al.*, 2007). The sulfated polysaccharides play a major role in treating inflammation. Also it was demonstrated that the fucans of *Fucus* species, *Laminarian* species on injection into rats reduced the peritoneal inflammation by leukocyte inhibition (Cumashi *et al.*, 2007). The ulvan polysaccharides of *Ulva rigida* also exerted anti inflammation by reducing immune stimulation (Jiao *et al.*, 2011). The fucoxanthin from seaweeds was active against inflammation and allergic reactions by degranulating the mast cells which secretes histamine (Shota Sakai *et al.*, 2009). It is noted that the carrageenan produced from *Euclima* or *Chondrus* or *Hypnea* species was used to analyse anti-inflammatory activity (Vipul *et al.*, 2014).

Applications of Seaweed

Macroalgal species are being used as a major viable source in various fields of applications. The carbohydrates content of seaweed has been the major area of interest in several fuel, food and medicinal industries. Many food additives like agar, carrageenan, laminarin, fucoidan extract from seaweed is found to be implemented in

food industries. Since the nanoparticles has gathered interest among researchers in the current world, seaweed nanoparticles being a natural source has potent application in biomedical industries apart from several human uses seaweeds are used as feed for cattle and aquatic animals and as fertilizer for various food and cash crops the various species of seaweed and its targeted content of application has been discussed in the following sections.

Biofuels

As there is a tremendous increase in fuel consumption these days, mankind has been in pursuit of natural source for fuel recovery which ensures pollution free environment. And now scientists have found certain seaweeds can be the raw material to produce fuel. These macroalgal biomass must be pre-treated for most biofuel applications. The first step of pre-treatment is to remove foreign objects and debris such as stones, sand, snails, or other litter that may be caught in the biomass either manually or by washing in many cases, chopping or milling is then required to increase the surface area/volume ratio (Fionnuala *et al.*, 2013). Finally, the biomass should be dewatered to 20%–30% to increase shelf life and reduce transportation costs in situations where it must be stored for long periods or transported over long distances before further processing.

The principal energy process considered for seaweed is fermentation, either anaerobic digestion, to create biogas, or ethanol fermentation. Other thermochemical options for macroalgae utilization include direct combustion, gasification, pyrolysis and liquefaction (Wei *et al.*, 2013). For better understanding about the production of biofuel from the algal biomass is made through the figure 5.

Bioethanol

Generally, bioethanol is produced from wood, grasses, and other inedible parts of plant but it is a tedious process to make sugar monomers. In order to overcome this, marine algae can be used as source for bioethanol production. Algae contain large quantities of carbohydrate biomass and high photon conversion efficiency for bioethanol production (Leilei Ge *et al.*, 2011, Hornl *et al.*, 2000). In addition, marine algae has buoyant property which can simplify the process of bioethanol production by neglecting the pre-treatment steps. It is noted that in Japan they use 4.47 million sq.km for harvesting *Sargassum horneri* for bioethanol production. *Ulvareticulata* macroalgae which can grow quickly was said to possess the potential to produce bioethanol. It is reported that other species such as *Padina japonica*, *Sphacelaria rigidula*, *Dictyosphaeria cavernosa*, *Sargassum polyphyllum* have appreciable dry weight content (Okazaki *et al.*, 1986). Brown seaweed *Laminaria hyperborea* has high amount of soluble carbohydrate (Ayhan Demirbas and M. Fatih Demirbas., 2011). Bioethanol from algae was known to reduce greenhouse gas emission by 85% over reformulated gasoline (Lalitesh *et al.*, 2014).

Biodiesel

Since diesel price is on hike irrespective of the demand biodiesel which is a natural fuel can be made from oils available within the algae. It is significantly noted that the macroalgae contains oil within its cell and biodiesel obtained with this oil is engine compatible. There are different varieties of macroalgae which contain oil once grown, the oil is removed from the macroalgae using chemicals or by squeezing oil out of the cells using scientific equipment.

Then the oil is used as an ingredient in biodiesel. This oil is changed chemically from plant oil to biodiesel. The finished product can be used on its own as pure biodiesel but is normally mixed with ordinary diesel and used directly in cars. Biodiesel is an alternative biodegradable energy source which has less CO₂ and no emission (Rocio Maceiras *et al.*, 2011). Oil extraction from algae and transesterification process are the two major steps involved in biodiesel production and the brown seaweed *Ascophyllum nodosum* has high oil content compared to the species such as *Sargassum*, *Codium*, *Ulva*, *Enteromorpha* (Van der Wal *et al.*, 2013).

Biogas

For use in domestic and industrial gas fuel seaweed can be subjected to anaerobic digestion for the production of methane gas it was estimated that the methane yield from anaerobic digestion of seaweed was 0.12 N/CH₄/g (Thomas Potts *et al.*, 2012). *Saccharina latissima* is a brown seaweed rich in carbohydrate was anaerobically digested for biogas production. It was reported that two process parameters such as steam explosion and thermal pretreatment method for biomass degradation seemed to affect the biogas yield in general the brown algae are more easily degraded than the green algae, and the green are more easily degraded than the red (Fionnuala Murphy *et al.*, 2013). Anaerobic digestion has been the most efficient method for the production of biogas rather than fermentation and thermal treatment. The biogas yield from certain species viz *Saccorhiza polyschides*, *Ulva species*, *Laminaria digitata*, *Fucus serratus* and *Saccharina latissima* are discussed in the following table 3 (Adam D Hughes *et al.*, 2012) despite the variations in the quantity and material ratios, the ratio between the methane produced and the input chemical

oxygen demand was reported to be stable (T. Matsui and Y. Koike., 2010).

Biobutanol

To note that the biobutanol has become a very competitive renewable biofuel for use in internal combustion engines and has been a boom to mankind. While comparing fuel properties it indicates that n-butanol is potent enough to remove the drawbacks brought by low-carbon alcohols or biodiesel. The applications of butanol as a biofuel are considered as three aspects, and they are as combustion experimentors in some well-defined burning reactors, as gasoline in spark ignition engine, as diesel fuel in compression ignition engine. From these demonstration that butanol is estimated effective as a second generation biofuel, from the viewpoints of combustion characteristics, engine performance, and exhaust emissions (Chao Jin *et al.*, 2011). *Ulva lactuca* was considered as the reserve species for acetone butanol ethanol (ABE) fermentation (Vivekanand *et al.*, 2012). The fermentation done with the bacterial strains such as *Clostridium beijerinckii*, *Clostridium saccharoperbutylacetoneum* and the algal carbohydrate resulted in the production of biobutanol. The yield and the concentration of biobutanol obtained from media were 0.29g butanol/g sugar and 4g/litre respectively (C. H. Vanegasa and J. Bartletta., 2013).

Polysaccharides of Seaweed

Seaweed polysaccharide is known for their varied functions and structures. They constitute natural sugars and sugar acids similar to land plants and in animals polysaccharides. As these polysaccharides contain hexose sugar, glucose, galactose and mannose, they have identical chemical formula, shape, properties and specific

atomic orientation (E. Hill and E. Surrey., 2007).

Agar

Seaweed agar is the most ancient phycocolloid found in Japan and discovered by Minoya tarozaemon in 1658 and first time manufactured in monument (Christopher S. Lobban, Michael J. Wynne., 1981, 1981). Agar is the major component of the cell-wall of certain red algae, which are the members of families, Gelidiaceae, Gelidiellaceae and Gracilariceae (D. Chakraborty., 1945). Agar consists of a chain of 9-p galactopyranose units linked in 1,4 bonds with a sulphated galactose in order to increase the yield and gel strength of agar, an alkaline treatment was done with sodium hydroxide for nearly one hour at the rate of 2 to 3 % alkali solution of 20,000 l/tonne at 90°C (Vigneswara Rao *et al.*, 1965). Also the sun-bleached seaweed was washed well in water and soaked for 24 h and then ground to a pulp and rinsed again in water and then the pulp was then extracted with water under pressure for 2 h after bringing the pH to 6 by adding of acetic acid. The agar gel was subjected to freeze thawing and bleached with NaClO before drying in a current of hot air (Taratra *et al.*, 2010). It is insoluble in cold water but soluble in boiling water. When agar was cooled to 34-43°C it forms a firm gel and does not melt further below 85°C (K. Funaki and Y. Kojima., 1951). In food technology agar is used as gelling and thickening agent in the confectionary and bakery industries, as stabilizer in the preparation of cheese and for salad dressings. In fish and meat industry, agar is applied for canned products, as a protective coating to avoid shaking during transport of these products. The agarose polycolloid play a prominent role in the DNA research and gel electrophoresis further agar is widely used in pharmaceutical industry as laxatives

as drug vehicle and as a medium for bacterial and fungal cultures and also used as an ion exchanger in the ion exchange resins (V. Rao and I.A.B. Thivy., 1960).

Alginate

Algin or alginic acid is a membrane mucilage and a major constituent of all alginates and the trade name is sodium alginate. Alginic acid is obtained from brown seaweed species such as Ecklonia, Macrocystis, Undaria, Laminaria and Duruilla from temperate area and turbinaria, sargassum, cystoseira and harmophysa from the tropical areas. Alginic acid which is the major polysaccharide of the brown seaweeds consists of unbranched chains comprising of contiguous β -1,4-linked D-mannuronic acid and blocks of contiguous α -1,4-linked L-guluronic acid (Arne Haug *et al.*, 1967). The proportions of D-mannuronic acid and L-guluronic acid varies between different species and from different parts of the same weed (A. Haug *et al.*, 1974). Alginates are found in both the intercellular region and the cell walls and it does biological functions structural and ion exchange type. Also it was extracted from the species laminaria digitata by alkaline extraction protocol (Peggy Vauchel *et al.*, 2008, Haug *et al.*, 1969). The molecular weight of alginate ranges generally between 500 and 1000 kDa (Moe *et al.*, 1995). As discussed earlier the alginate composition of different seaweed species are ascophylum nodosum (22–30%); laminaria digitata fronds (25–44%); laminaria digitata stipes (35–47%); laminaria hyperborea fronds (17–33%); laminaria hyperborean stipes (25–38%) (Rinaudo M., 2007.). Alginate contents ranges between 17 and 45% are extracted in sargassum species (E. Fourest and B. Volesky., 1996). In industries the alginates are extracted from brown seaweed sargassum turbinarioides by cutting the

thallus with a knife near the rhizoid and the algae were washed and sun-dried at ambient temperature and stored in aerated bags (Taratra *et al.*, 2010). Japanese work on the brown seaweed has revealed that alginate is subjected to polymerization in the cytoplasm and then transported to the cell surface (Abe *et al.*, 1973). It has observed d-mannuronic acid precursor of polymannuronic acid that while separating from the brown seaweed, *Fucus gardneri* in young tissue. They identified the presence of trace quantity of a gdp-guluronic acid (T.Y. Lin and W.Z. Hassid., 1966). Further on extracting epimerase from *pelvetia canaliculata* the conversion of polymannuronic acid into a mixed poly-d-mannuronic-l-guluronic polymer has been witnessed by tritium incorporation (Madgwick *et al.*, 1973). In beverages alginates acts as clarifying agents for making wines and liquor where as it acts as foam stabilizer in lager beer and malt beer (H. Ertesvåg and S. Valla., 1998.). Artificial casings are made with alginates as poses to ensure longer shelf life in sausage industries and it alginates are used in the form of gel for deep freezing of fish, meat and poultry products in western countries (Bernd H. A. Rehm. 2009).

Carrageenan

Macroalgal carrageenan is a sulphated polymer obtained from various red seaweeds and it differs from agar in its high sulphate and ash content. The major difference between the agars and carrageenan is that the former contains d- and l-galactose units whereas the latter consists entirely of the d-sugar (E. Hill and E. Surrey., 2007). They are commercially important hydrophilic colloids present in the matrix of red seaweeds (rhodophyta) and does structural function also they are considered as high sulfated galactans and as strong anionic polymers. Seaweed species *Kappaphycus*, *Eucheuma*

Chondrus crispus, *Gigartina stellata*, *Iridaea*, *Hypnea* species have high content of Carrageenan (H.J. Bixler and H. Porse., 2011). Carrageenan can be recovered to either by direct drying on steam-heated rolls or by precipitation of the carrageenan from solution by 2-propanol or other alcohols. It is to be noted that in the past alcohol precipitation method was used to recover carrageenan from irish moss (G. Bourgade., 1871).

Fucoidan

Fucoidan is a sulfated polysaccharide found mainly in various species of brown seaweed and fucoidan is used as dietary supplements (M. Tutor-Ale *et al.*, 2011). Fucoidan is a class of sulfated, fucose rich, polysaccharides found in the fibrillar cellwalls and intercellular spaces of brown seaweeds. Fucose-containing sulfated polysaccharides (fcsp) consist of a backbone of (1→3)- and (1→4)-linked α -l-fucopyranose residues, that may be organized in stretches of (1→3)- α -fucan or of alternating α (1→3)- and α (1→4)-bonded l-fucopyranose residues (M. Tutor-Ale *et al.*, 2011). Apart from fucose and sulfate groups fucoidans also contain galactose, xylose, mannose and other uronic acid (M. Tutor-Ale *et al.*, 2011). Fucoidan was extracted using dilute acetic acid from various species of *Laminaria* and *Fucus* [201]. Fucoidans were reported to possess various biological such as anti-inflammatory, anticoagulant, antithrombotic (Trincherro *et al.*, 2009, Raghavendran *et al.*, 2011), antiviral including anti-HIV (Kusaykin *et al.*, 2008, Lee *et al.*, 2004), immunomodulatory (Raghavendran *et al.*, 2011), antioxidant (Wang *et al.*, 2010), and antitumor (Andriy *et al.*, 2010). Fucoidan from *Laminaria* species were found to inhibit a variety of DNA and RNA enveloped viruses and also useful in elucidation of

mammalian sperm and egg (Frank *et al.*, 1989). The "fucan" extracted from *Pelvetia canaliculata* had a very strong affinity for magnesium which in turn can assist the contact of their fronds with seawater. It was also studied that *Pelvetia canaliculata*, since grows on the higher part of the shores have a high "fucan" content (Carlberg *et al.*, 1978).

Laminarin

Laminarin is a storage glucan found in brown algae and is used as a carbohydrate food reserve similar to diatoms (Anne Beattie *et al.*, 1961). Laminarin is β -glucan and it is known to induce anti-apoptotic and anti-tumoral activities (Rioux *et al.*, 2010). This is a water-soluble polysaccharide containing approximately 20-25 glucose units. It has two types of chains namely, g-chains terminated at the reducing end with glucose and m-chains terminated by mannitol (Percival E., 1978). In addition the laminaran, from *Eisenia bicyclis*, may contain 1,6-linked units in the chains, or the chains may be branched at C-6 (Bidwell, 1967). It was that the mannitol and laminarin are active metabolites which can be interconverted (Quatrano RS and Stevens PT., 1976). Also it was shown that laminarin isolated from the cytoplasm of developing zygotes of *Fucus* species decreased during the first 7 hours of wall assembly while the content in cellulose in the wall increased laminarin structure and composition vary according to algae species (Chizhov *et al.*, 1998). Based on the degree of polymerization the molecular weight of laminaran has been found as 5000KDa approximately (Rioux *et al.*, 2007).

Pharmaceutical Application of Seaweed

Most inspired bioresource as far as pharmaceuticals is concerned is seaweed because of the high nutrient content.

Seaweed has been used as food throughout Asia, Japan, China, Rome, etc., to treat various health disorders. The romans used seaweed in the treatment of wounds, burns, and rashes (Hocman G., 1989) whereas in Scotland physicians used dried seaweed stem to drain abdominal wall abscesses and they also inserted seaweed into the cervix to treat dysmenorrhea and seaweed was also employed intravaginally for vaginal atresia for which it was used urethrally and rectally for strictures (Abdussalam S., 1990, Johnson N., 1990). Traditional Chinese medicine includes use of the brown alga *Laminaria* in the treatment of cancer. The ancient Egyptians used seaweed to treat breast cancer. Seaweeds are being extensively used in cardiovascular conditions as it can reduce cholesterol level. In general alginates from seaweeds has been used in wound dressings and as fillers in tablets, pills and as ointment base whereas seaweed carrageenan acts as good emulsifiers in mineral oil and drug preparations. *Saccharina japonica* and *Undaria pinnatifida* was analysed to contain fucoidan which can destruct cancer cells. Also complete wipe out of cancer cells by fucoidan treatment has been demonstrated in Japan. It is noted that cancer mortality rates and breast cancer rates are considerably low in Japan because of seaweed consumption (Jane Teas *et al.*, 2009, Carmen Aceves *et al.*, 2005). Seaweed extracts being a source of calcium, magnesium, selenium and other minerals has been evaluated to treat osteoarthritis (Frestedt *et al.*, 2009, Joy *et al.*, 2008). And it was reported that intake of seaweed powder 5g/day, 12g/day and 4-6g/day in diet controls cholesterol, hypertension, metabolic syndrome respectively (Yukio *et al.*, 2001, Krotkiewski *et al.*, 1991). Iodine rich seaweeds like *Asparagopsis tucjirmis*, *Sarcocystis* species was reported to cure hypothyroidism (goitre). Also in the earlier

research works seaweed extracts was known to stimulate B lymphocytes and macrophages that in turn modulates immune response (Liu *et al.*, 1997, Shan *et al.*, 1999).

Application of seaweed in other areas

On pursuing the trends of biotechnology and its advancements, seaweed a single macroalgal source with varied types, characteristics and multipotent properties was known to be employed in various sectors as from animal feed to the latest emerging field nanotechnology. From the earlier literature works done the other areas of effective uses of seaweeds and their distinct types are summarised in the following sections.

Seaweed Nanoparticles

Increasing trend in nanotechnology has gathered interest among the researcher thereby algal nano particles are known as bio nano factories as they are highly stable, easy to handle and avoids cell maintenance. Added to this, metal nano particles from seaweed have excellent potential in biomedical applications (Song JY and Kim BS., 2009). Benign nanoparticle synthesis which is nontoxic has been an emerging trend in today's world (Shankar *et al.*, 2004). Ecofriendly gold nanoparticles synthesized from *Turbinaria conoides* was confirmed to be associated with carboxylic, amine, and polyphenolic groups by fourier transform-infrared spectroscopy (Shanmugam *et al.*, 2013). Green seaweed *Caulerpa peltata*, red *Hypnea valencia*, brown *Sargassum myriocystum* seaweeds were used to synthesize zinc oxide nanoparticles and they can be used in effluent treatment process to reduce microbes (S. Nagarajan and K.A. Kuppusamy., 2013). Silver nanoparticles from *Sargassum tenerrimum* along with the

presence of phytochemicals as reducing agents was found to have excellent antimicrobial activity (Dhanalakshmi *et al.*, 2012). Further report identified that Fe_3O_4 nanoparticles obtained from *Sargassum muticum* which contain sulfated polysaccharides as the reducing agent was determined to have antimicrobial potential, stabilizing capacity and this on fabrication came out with other metal oxides (Mahnaz *et al.*, 2013).

Seaweed in Wastewater Treatment

The two major areas of waste water treatment in which seaweed has its prominent role to play are one is to treat sewage and agricultural wastes to exploit nitrogen- phosphorous wastes and the other is to remove toxic metals from industrial effluent and it was suggested to be a boon in coastal areas (Schramm *et al.*, 1991). Alginates extracted from the seaweed *Sargassum sinicola* was used to co-immobilize the microalgae *Chlorella sorokiniana* and the bacterium *Azospirillum brasilense* which promotes growth when employed in waste water treatment. Seaweeds were used in wastewater treatment because of its ability to absorb nutrients and heavy metal ions that are toxic. Also seaweeds were suggested as biological indicator of marine pollution like eutrophication on considering its capacity to take up ammonia in the nitrogen form and phosphorous. Despite varying concentration and type seaweeds such as *Sargassum*, *Laminaria*, *Ecklonia*, *Ulva* and *Enteromorpha* were identified as indicators of heavy metal pollution (Ryther *et al.*, 1975).

Seaweed as beauty promoters

Seaweeds possess potentials to stimulate blood circulation and revitalize, nourish and eliminate toxins of the skin. Seaweed bath is

one such instance been practiced in Ireland with fucus serratus species promoted to treat rheumatism and arthritis. Also all the required nutrients, aminoacids and oils were found to be absorbed by the skin in seaweed bath. An Irish company has been established to produce seaweed powder from Ascophyllum nodosum for the cosmetic and algotherapy and it was said to improve damaged hair by means of ionic interactions

with the proteins of hair. Also seaweeds enzymes were found to heal dandruff and stimulate hair follicles for hair growth. Silicon from seaweeds were analysed to have anti- wrinkle effect on facial skin and other anticellulite preparations from seaweeds has been used in the form of creams and lotions for hip, thigh and neck (Pramanick *et al.*, 2013).

Table.1 Biochemical composition of some seaweed species

Species	Biochemical compound	Yield
<i>Gracillariaspecies</i>	vitamin A	865µg RE /100g
	vitamin C	28.5± 0.1mg/100g
	Fiber	(24.7±0.7)% wet wt
	Lipid	0.43-(3.3±0.2)% wet wt
	PUFA	2.43-51.77% w/w
	Protein	(6.9±0.1)% wet wt
	Minerals	153.3±2.64mg/100g
	Ash	(22.7 ±0.6)%
	Aminoacid	116.37mg/g
<i>Padinaspecies</i>	Carbohydrate	50.9±0.52mg/g dry wt
	Lipid	58.38-184.0 mg/g dry wt
	Protein	17.1% d.wt
	Vit B6	115±0.577 µg/g
	Vit B2	75±1.732µg/g
	Vit B1	46±2.309µg/g
	Minerals	90.93±0.257 µg/g
	Trace elements	30-110 ppm
<i>Enteromorphaspecies</i>	Omega -3 FA	10.38%
	Omega -6FA	10.9g/100g
	Fiber	33.4%dry wt
	Protein	19.5%
	Carbohydrate	64.9%
	Ash	15.2%
	Lipid	0.3%
	Moisture	9g/kg

<i>Ulva</i> species	Carbohydrates	(17.8±2.3)-70.1% d.wt
	Protein	14.7%
	Minerals	247.54mg/100g d.wt
	Lipids	0.51-1.18 %
	SFA	65.40±1.65%
	Aminoacids	829.48±6.51mg/100g
	Fibre	51.3-62.2% d.wt
	Ash	25.9-62.2% d.wt
	Vitamin E	19.70 µg/g d.wt
<i>Laminaria</i> species	Carbohydrates	23.61%
	Protein	16±3.8g/100g
	Lipids	26.05±0.3g/10g
	Aminoacids	86.5±14.1g/100gN
	Fibre	36.0±5.7g/100g
	Ash	24.9-36.4%
	Minerals	1929.57mg/100g d.wt
	Vitamin E	34.38 µg/g d.wt
<i>Porphyra</i> species	Carbohydrates	45.1%
	Protein	47% d.wt
	Lipids	2.1±1.2g/100g
	Fibre	47.7±4.25g/100g
	Ash	20.59±0.16g/100g d.wt
	Minerals	106.07±3.44
<i>Palmaria</i> species	Protein	28.25% d.wt
	Minerals	1686.78mg/100gd.wt
	Fatty acids	25.53±2.37%
	Moisture	84%
	Ash	12-37%
<i>Undaria</i> species	Ash	39.26±0.24g/100g d.wt
	Minerals	9992.42±383mg/100g d.wt
	Fibre	35.3%
	Fatty acids	33.36±3.91g/100g d.wt
<i>Sargassum</i> species	Carbohydrate	29.37-47.04±0.7% d.wt
	Protein	11.52-16.59±0.86%
	Fattyacid	0.16±0.006-20.96%
	Fibre	77.40±16.14g/kg
	Ash	194.30±48.65g/kg
	Mineral	1901.74mg/100g d.wt
	Moisture	14.33±1g/kg

<i>Fucus</i> species	Carbohydrate	65.7±0.4%
	Lipid	1.4±0.1%
	Protein	7%
	Mineral	11736±127mg/100g d.wt
	Ash	0.11-22.71±3.1%
	Vitamin	820 µg/gd.wt
<i>Hypnea</i> species	Carbohydrate	13.69%
	Protein	28.63±0.39%
	Lipid	1.43%
	Fibre	47.0025%
	Ash	21.28%
	Moisture	11.18%
	Aminoacid	77.1g/100g
<i>Gelidium</i> species	Protein	11.55±1.02%
	Carbohydrate	41.87±2.21%
	Lipid	1.53±0.61%
	Fibre	24.74±1.05%
	Ash	17.57±0.74%
	Moisture	10.85±0.98%
<i>Turbinaria</i> species	Protein	14.68%
	Carbohydrate	18.46±0.685%
	Lipid	2.53±0.14%
	Moisture	83.79±0.32%
	Ash	21.37±0.60%
	Fibre	29.61±1.59% d.wt
<i>Acanthopora</i> species	Carbohydrate	23.54±0.10%
	Lipid	55.33±5.1g/mg
	Fattyacid	64%
	Ash	16.4±2.4g/mg
	Protein	24%
<i>Colponemias</i> species	Carbohydrate	32.1±1.75%
	Protein	9.2±1.78%
	Lipid	1.5±0.29%
	Moisture	11.5±0.20%
	Fatty acid	1.0%
	Ash	28.1±0.86%
<i>Ascophyllum</i> species	Carbohydrate	69.6±0.2%
	Protein	(3-15)% d.wt
	Lipid	1.2±0.1 %
	Oil	2.85 %
	Ash	22.5±0.1 %
	Vitamin C	81.75 µg/g d.wt.
	Vitamin E	3.63 µg/g d.wt

<i>Dictyota</i> species	Carbohydrate	10.63%
	Lipid	16.1-20.2 % d.wt
	Protein	9.8%
	Minerals	0.559±0.009 ppm
<i>Caulerpa</i> species	Carbohydrate	9.7±0.477 %
	Protein	24.55±0.84 %
	Lipid	0.9±0.38%
	Fiber	1.36 g/kg
	Ash	24.2 % d.wt
<i>Jania</i> species	Protein	2.53±0.15% wet wt
	Lipid	0.22±0.035% wet wt
	Moisture	40.13±1.54 % wet wt
	Ash	48.14±1.88 % wet wt
	Fatty acid	11.319 % wet wt

Table.2 Biogas yield from certain seaweed species [16]

Species Name	Yield in ml g volatile solids(g_{vs}^{-1})
<i>Saccharinalatis sima</i>	335
<i>Saccorhiza polyschides</i>	255
<i>Laminaria digitata</i>	246
<i>Ulva sp.</i>	191

Fig.1 General structure of a seaweed




Fig.2 Distribution of species along Indian Ocean




Fig.3 Graphical representation of species distribution




Fig.4 Macroalgal species weightage in other countries





Fig.5 Flowchart showing various biofuel production from algal biomass



Seaweed as Animal Feed

Seaweed meal is rich in proteins and carbohydrates has been used as animal feed after getting the raw biomass cleaned, dried and milled. Also seaweeds were reported to be used as domestic animal feed in Norway, Iceland and France for years. Seaweed feed is considered to have 30 percent of the nutrition of the grains. It was being added to poultry diets in a ratio of 5-15 percent of the diet depending upon seaweed species and the targeted animal age. In addition feeding seaweed meal and sardine oil to chickens has resulted in the reduction of cholesterol level and increase in omega -3 fatty acid content. Seaweed extract from *Ascophyllum nodosum* on consumption by pigs was known to promote prebiotic activity as it has ascorbic acid, tocopherols, vitamin B, minerals and it was found to provide iodine for hogs, chickens, sheep and dairy cows (Jensen, A., 1972). The wet seaweed were being processed by passing through hammer mills with small screens to get fine particles followed by passing through drum dryer at 700-800°C. It was reported that *Ascophyllum* which is a dark brown seaweed has high content of phenolic compounds whereas *Alaria esculenta* which is light brown seaweed was found to be more effective as animal feed than the latter. *Macrocystis pyrifera* and *Sargassum* species was evaluated as fodder supplements in goat and sheep diets. Also *Sargassum* species was fed to Nubian goats which replaced 25% flour feed (M Casas-Valdez and E.A. Maina., 2006).

Seaweed as a Biofertilizer

Seaweeds have been used to nourish worn out soil around coastal areas. Use of seaweed fertilizer stimulated root volume, plant growth and even promoted fruit development thus resulted in the

production of high quality agricultural products (Pramanick *et al.*, 2013). Further it was revealed that the use of seaweed fertilizer improved germination and disease resistant capacity in plants (Khan *et al.*, 2009). Water and alkaline extracts of *Ascophyllum nodosum* has given tomatoes of appreciable mass and also yielded good quality fruits. Seaweed extracts and fertilizers have given promising turnouts in agriculture for which it is a main stay in the catalogue of commonly preferred soil nourishers (M. Povolny, 1981).

Indian scenario

In the earlier works it has been reported that there are 271 genus and 1153 species of marine algae found along the Indian coast. It was estimated that the southern coast of India has got more than 200 species of seaweeds which stands a prominent source for agar and sodium alginate production in India. Also it was found that 75,373 tonnes of seaweeds the maximum mass are scattered along the Tamilnadu coastal line covering an area of 1863 sq km from Rameshwaram to Kanyakumari. It was reported that there are about 25 seaweed processing small scale industries of which 20 are agar producers however 110-132 tonnes of dry agar are being produced annually based on the harvest of 880-1,100 tonnes (dry weight) in Tamilnadu and Kerala. Seaweeds like *Gracilaria*, *Gelidium*, *Kappaphycus* are being cultivated on a large scale in India for the production of phycocolloids (Ayhan Demirbas and M. Fatih Demirbas., 2011). Though India has got a longer coastal line and rich availability of seaweed resource the awareness in India about seaweed culture and its commercial applications is not fair enough. So the common people need more awareness about seaweeds to promote a better healthy future.

Economic Importance

From all the seaweed potentials studied earlier it is evident that the globalization of seaweed cultivation and seaweed derived products can improve the economic status of various countries all over the world. The European, Eastern and Southeast Asian countries are the major producers and consumers of seaweeds (Murty U.S and Banerjee A.K. 2012). According to FAO 2014 Chile tops the global natural seaweed producers and it was estimated as one of the first agar producers in the world (H.J. Bixler and H. Porse., 2011). Various seaweeds are being harvested for human consumption and industrially used in the production of a wide range of nutraceuticals. Patagonia is an outstanding place where all the seaweeds are harvested and commercialized as such biomass and bioproducts (Céline *et al.*, 2014). The red seaweeds are harvested for agar and carrageenan industries whereas the brown species are utilized by alginates industries (Zaixso *et al.*, 2006). It is to be noted that rapid growth is seen in Japan and Korea is mainly because of seaweed production. The 2010 estimate has shown that the global seaweed production was 19 million tons and it accounts for US \$5.7 billion. The increasing demand for seaweed based edible products puts its cultivation in a profiting trend.

In conclusion, seaweed thus serves as a sustainable feedstock and an ecofriendly resource for various purposes. Also many bioactive compounds and pharmacologically active substances have been isolated from macroalgae and put into use in various forms. Nowadays seaweeds are under threat in developing nations because of human settlements, natural barriers and lack of awareness regarding its potential use. Awareness about the importance of harvesting and commercialization of

seaweeds and products should be effectively promoted in developing nations. The cultivators should be well trained to make them familiar with various harvesting techniques, the parameters and specificity involved in growing various classes of species for successful commercialization of seaweed farming. India, since it has got long coastal line with rich availability of seaweed species, if encouraged and promoted with financial aids by the government, malnutrition and poverty can be uprooted to the maximum extent.

References

- Haug, B. Larsen, B Smidsrod., 1974. Uronic acid sequence in alginate from different sources, *Carbohydr. Res.* 32, (2), 217–225.
- Haug, B. Larsen, E. Baardseth., 1969. Comparison of the constitution of alginates from different sources. *Proc. int. Seaweed Symp* 6, 443–451.
- A.B. Zaixso, M. Ciancia, A.S. Cerezo. 2006. Seaweed resources in Argentina. World seaweed resources—an authoritative reference system, Margraf publishers GMBH (2006), Germany.
- Abdussalam S., 1990. Drugs from seaweeds. *Med Hypotheses.* 32(1), 33-35.
- Adam D Hughes, Maeve S Kelly, Kenneth D Black, Michele S Stanley., 2012. Biogas from Macroalgae: Is it time to revisit the idea?, *Biotechnology for Biofuels*, 5, (1), 86. doi:10.1186/1754-6834-5-86.
- Aleksander Handa, Silje Forbord, Xinxin Wang, Ole Jacob Broch, Stine Wiborg Dahle, Trond Røvik Størseth, Kjell Inge Reitan, Yngvar Olsen Jorunn Skjermo *et al.*, 2013. Seasonal- and depth-dependent growth of cultivated kelp (*Saccharina latissima*) in close proximity to salmon (*Salmo salar*) aquaculture in Norway. *Aquaculture*. 414, 191–201.
- Alekseyenko TV, Zhanayeva SY, Venediktova AA, Zvyagintseva TN, Kuznetsova TA, Besednova NN, Korolenko TA., 2007. Antitumor and antimetastatic activity of fucoidan, a sulfated polysaccharide isolated from the okhotsk sea fucus evanescens brown alga. *Bulletin of experimental*

- biology and medicine, 143(6) 730-732.
- Andriy Synytsya, Woo-Jung Kim, Sung-Min Kim, Radek Pohl, Alla Synytsya, František Kvasnička, Jana Čopíková, Yong Il Park., 2010. Structure and antitumour activity of fucoidan isolated from sporophyll of Korean brown seaweed *Undaria pinnatifida*, *Carbohydr. Polym.* 81 (1), 41–48.
- Anne Beattie, E. L. Hirst, and Elizabeth Percival., 1961. Studies on the metabolism of the Chrysophyceae. Comparative structural investigations on leucosin (chrysolaminarin) separated from diatoms and laminarin from the brown algae. *Biochem. J.* 79(3), 531–537.
- Arne Haug, Bjørn Larsen, Olav Smidsrød, G. Eriksson, R. Blinc, S. Paušak., 1967. Studies on the sequence of uronic acid residues in alginic acid, *Acta Chemica Scand.* 21, 691–704.
- Ayhan Demirbas, M. Fatih Demirbas., 2011. Importance of algae oil as a source of biodiesel, *Energy Conversion and Management.* 52(1), 163–170.
- Bansemir A., Blume M., Schröder S., Lindequist U., 2006. Screening of cultivated seaweeds for antibacterial activity against fish pathogenic bacteria, *Aquaculture.* 252, 79–84.
- Bartolomeu W.S. Souza ., Miguel A. Cerqueira., Ana I. Bourbon., Ana C. Pinheiro., Joana T. Martins., José A. Teixeira., Manuel A. Coimbra., António A. Vicente., 2012. Chemical characterization and antioxidant activity of sulfated polysaccharide from the red seaweed *Gracilariabiridae*, *Food hydrocolloids.* 27, 287–292
- Bernd H. A. Rehm. 2009. *Alginates: Biology and Applications* Springer Dordrecht Heidelberg London New York DOI: 10.1007/978-3-540-92679-5.
- Bidwell, R. G. S., 1967. Photosynthesis and metabolism in marine algae. VII. Products of synthesis in fronds of *Fucus vesiculosus* and their use in respiration. *Can. J. Bot.*, 45: 1557–1565.
- Brian E. Lapointe, F. Xavier Niell and Jose Miguel Fuentes., 1981. Community structure, succession, and production of seaweeds associated with mussel-rafts in the Ria de Arosa, N.W. Spain. *Marine Ecology - Progress Series* 5, 243–253.
- Buschmann, A. H., Westermeier, R., Retamales, C., 1995. Cultivation of *Gracilaria* in the sea-bottom in southern Chile: a review. *J. Appl. Phycol.* 7 (3), 291–301.
- A. H. Vanegasa and J. Bartletta., 2013. Green energy from marine algae: biogas production and composition from the anaerobic digestion of Irish seaweed species, *Environmental Technology*, 34, (15), 2277–2283.
- C.L.F. De almeida., H.D.S. Falcao., G.R. Dem.lima., 2011. Bioactivities from marine algae of the genus *gracilaria*, *Int j . mol.sci.* 12 (7), 4550–4573.
- Carmen Aceves, Brenda Anguiano and Guadalupe Delgado., 2005. Is iodine a gatekeeper of the integrity of the mammary gland?, *J Mammary Gland Biol Neoplasia.* 10 (2), 189–196.
- Catherine Vonthron-Sénécheau, Marcel Kaiser, Isabelle Devambe, Antoine Vastel, Isabelle Mussio and Anne-Marie Rusig., 2011. Antiprotozoal activities of organic extracts from french marine seaweeds, *Marine drugs*, 9(6), 922–933.
- Céline Rebours, Eliane Marinho-Soriano, José A. Zertuche-González, Leila Hayashi, Julio A. Vásquez, Paul Kradolfer, Gonzalo Soriano, Raul Ugarte, Maria Helena Abreu, Ingrid Bay-Larsen, Grete Hovelsrud, Rolf Rødven, and Daniel Robledo. 2014. Seaweeds: an opportunity for wealth and sustainable livelihood for coastal communities. *J Appl Phycol.* 26(5), 1939–1951.
- Celum CI., 2004. The interaction between herpes simplex virus and human immunodeficiency virus. *Herpes.*; 11: 36A–45A
- Chao Jin, Mingfa Yao, Haifeng Liu, Chia-fon F. Lee, Jing Ji., 2011. Progress in the production and application of n-butanol as a biofuel, *Renewable and Sustainable Energy Reviews*, 15, (8), 4080–4106.
- Chizhov A.O, A. Dell, H.R. Morris, A.J. Reason, S.M. Haslam, R. A. McDowell, O.S. Chizhov and A.I. Usov., 1998. Structural analysis of laminarin by MALDI and FAB mass spectrometry, *Carbohydrate Research.* 310, 203–210.
- Christopher S. Lobban, Michael J. Wynne., 1981. *The Biology of Seaweeds*; ISBN: 9780520045859
- Cumashi A, Ushakova NA, Preobrazhenskaya ME, D'Incecco A, Piccoli A, Totani L, Tinari N, Morozevich GE, Berman AE,

- Bilan MI, Usov AI, Ustyuzhanina NE, Grachev AA, Sanderson CJ, Kelly M, Rabinovich GA, Iacobelli S, Nifantiev NE., 2007. A comparative study of the anti-inflammatory, anticoagulant, antiangiogenic, and antiadhesive activities of nine different fucoidans from brown seaweeds. *Glycobiology*. 17(5), 541-552
- Chakraborty. 1945. Agar-agar manufacture from *Gracilaria confervoides*, *Jour. Proc. Inst. Chem.* 17, 188.
- D.J. Newman., G.M.Cragg., K.M. Snader., 2003. Natural products as sources of new drugs over the period 1981-2002. *Journal of natural products* 66(7), 1022-1037.
- Dadollahi-Sohrab, A., Garavand-Karimi, M., Riahi, H., Pashazanoosi, H., 2012. Seasonal variations in biomass and species composition of seaweeds along the northern coasts of Persian Gulf (Bushehr Province). *J. Earth Syst. Sci.* 121(1), 241-250.
- De Souza, É.T.; Pereira de Lira, D.; Cavalcanti de Queiroz, A.; Costa da Silva, D.J.; Bezerra de Aquino, A.; Campessato Mella, E.A.; Prates Lorenzo, V.; De Miranda, G.E.C.; De Araújo-Júnior, J.X.; De Oliveira Chaves, M.C.; Barbosa-Filho, J.M.; Filgueiras de Athayde-Filho, P.; De Oliveira Santos, B.V.; Alexandre-Moreira, M.S., 2009. The Antinociceptive and Anti-Inflammatory Activities of Caulerpin, a Bisindole Alkaloid Isolated from Seaweeds of the Genus *Caulerpa*. *Mar. Drugs*, 7(4), 689-704.
- Dhanalakshmi. P. K., Riyazulla Azeez, Rekha. R, Poonkodi. S, Thangaraju Nallamuthu., 2012. Synthesis of silver nanoparticles using green and brown seaweeds, *Phykos* 42 (2) 39-45.
- E. Fourest and B. Volesky., 1996. Contribution of sulfonate groups and alginate to heavy metal biosorption by the dry biomass of *Sargassum fluitans*, *Environmental Science and Technology*, 30, (1), 277-282.
- [173]Peggy Vauchel, Raymond Kaas, Abdellah Arhaliass, Régis Baron and Jack Legrand., 2008., A New Process for Extracting Alginates from *Laminaria digitata*: Reactive Extrusion, *Food and Bioprocess Technology*, 1, (3), 297-300
- E. Hill and E. Surrey., 2007. The polysaccharides of green, red and brown seaweeds: Their basic structure, Biosynthesis and function. *Food and Bioprocess Technology*, 1, 297-300.
- Erwan Plouguerné, Lauro M. de Souza, Guilherme L. Sasaki, Jéssica Figueiredo Cavalcanti, Maria Teresa Villela Romanos, Bernardo A. P. da Gama, Renato Crespo Pereira and Eliana Barreto-Bergter., 2013. Antiviral Sulfoquinovosyl diacylglycerols (SQDGs) from the Brazilian Brown Seaweed *Sargassum vulgare*. 11(11), 4628-4640
- Farideh Namvar, Rosfarizan Mohamad, Javad Baharara, Saeedeh Zafar-Balanejad, Fahimeh Fargahi, and Heshu Sulaiman Rahman., 2013. Antioxidant, Antiproliferative, and Antiangiogenesis Effects of Polyphenol-Rich Seaweed (*Sargassum muticum*), *BioMed Research International*. 2013, 604-787.
- Fionnuala Murphy, Ger Devlin, Rory Deverelland Kevin McDonnell., 2013. Biofuel Production in Ireland-An Approach to 2020 Targets with a Focus on Algal Biomass, *Energies*, 6,(12), 6391-6412.
- Frank C. Church\$, James B. Meade, Rita E. Treanor, and Herber Ct . Whinna., 1989. Antithrombin activity of function. The interaction of fucoidan with heparin cofactor II antithrombin III, nad thrombin, *J. Biol-chem* 264, (6), 3618-3625.
- Frestedt JL, Kuskowski MA, Zenk JL., 2009. A natural seaweed derived mineral supplement (Aquamin F) for knee osteoarthritis: a randomised, placebo controlled pilot study, *Nutr J.* 8(7).
- Fu-Jin Wang., Sheen-Kye Kang., Chorong-Liang Pan., Shao-Chi Wu., 2011. Composition of two seaweed-oligosaccharide-lysates derived from agarase as-ii and their ferrous-chelating antioxidant power. *Journal of marine science and technology.* 19 (5)557-564
- G. Bourgade., 1871. Improvement in treating marine plants to obtain gelatine. *Journal of Applied Phycology*, 112,535.
- G.E. Carlberg, E. Percival, M.A. Rhaman., 1978. Carbohydrates of the seaweeds, *Desmarestia ligulata* and *D. firma.*, *Phytochem.* 17, (8), 1289-1292.
- Ghislain Moussavou, Dong Hoon Kwak, Brice Wilfried Obiang-Obonou, Cyr Abel Ogandaga Maranguy, Sylvatrie-Danne Dinzouna-Boutamba, Dae Hoon Lee, Ordélia Gwenaelle Manvoudou

- Pissibanganga, Kisung Ko, Jae In Seo and Young Kug Choo., 2014. Anticancer effects of different seaweeds on human colon and breast cancers, *Marine drugs* 12 (9) 4898-4911.
- Gupta, S. and N. Abu-Ghannam. 2011. Recent developments in the application of seaweeds or seaweed extracts as a means for enhancing the safety and quality attributes of foods. *Innovative Food Science and Emerging Technologies* 12(4): 600–609.
- H. Ertesvåg and S. Valla., 1998. Biosynthesis and applications of alginates, *Polymer degradation stability*, 59 (1-3) 85–91.
- H. Indu., R. Seenivasan., 2013. Invitro antioxidant activity of selected seaweeds from southeast coast of India. *International Journal of Pharmacy and Pharmaceutical Sciences*. 5 (2), 474-484.
- H.J. Bixler and H. Porse., 2011. A decade of change in the seaweed hydrocolloids industry, *Journal of Applied Phycology*, 23 (3) 321-335.
- H.J. Bixler, H. Porse., 2011. A decade of change in the seaweed hydrocolloids Industry, *J appl phycol*. 23, 321–335.
- H.R. Raghavendran, P. Srinivasan, S. Rekha, 2011. Immunomodulatory activity of fucoidan against aspirin-induced gastric mucosal damage in rats, *Int. Immunopharmacol.* 11, (2), 157–163.
- Harada H, Yamashita U, Kurihara H, Fukushi E, Kawabata J, Kamei Y., 2002. Antitumor activity of palmitic acid found as a selective cytotoxic substance in a marine red alga. *Anticancer res.* 22 (5) 2587–2590.
- Hasegawa, Y., 1971. Forced cultivation of *Laminaria*. In *Proc. Seventh Internatl. Seaweed Symp.*, Sapporo, Japan. 391-393.
- Hee-Kyoung Park, In-Hye Kim, Joongkyun Kim and Taek-Jeong Nam., 2013. Induction of apoptosis and regulation of erbB signaling by laminarin in ht-29 human colon cancer cells, *Int. j. mol. med.* 32 (2) 291–295.
- Hee-Kyoung Park, In-Hye Kim, Joongkyun Kim, and Taek-Jeong Nam., 2012. Induction of apoptosis by laminarin, regulating the insulin-like growth factor-ir signaling pathways in ht-29 human colon cells, *Int. J. Mol. Med.* 30 (4) 734–738.
- Hocman G., 1989. Prevention of cancer: vegetables and plants, *Comp Biochem Physiol B.* 93(2), 201-212.
- Immanuel G., Vincybai V.C., Sivaram V., Palavesam A., Marian M.P., 2004. Effect of butanolic extracts from terrestrial herbs and seaweeds on the survival, growth and pathogen (*Vibrio parahaemolyticus*) load on shrimp *Penaeus indicus* juveniles, *Aquaculture*. 236 (1-4) 53–65.
- Ismail A Jr., Tan S., 2002. Antioxidant activity of selected commercial seaweed, *Mal j nutr.* 8(2), 167-177.
- J. Madgwick, A. Haug, B. Larsen., 1973. Polymannuronic acid 5-epimerase from the marine alga *Pelvetia canaliculata* (Dcne et Thur.), *Acta Chem. Scand.* 27 (9) 3592–3594.
- J.H. Ryther, J.C. Goldman, C.E. Gifford, J.C. Huguenin, A.S. Wing, J.P. Clarnier, L.D. Williams, B.E. Lapointe., 1975. Physical models for integrated waste recycling-Marine polyculture system. *Aquaculture*. 5, 163-177.
- Jane Teas, Thomas G. Hurley, James R. Hebert, Adrian A. Franke, Daniel W. Sepkovic, and Mindy S. Kurzer., 2009. Dietary seaweed modifies estrogen and phytoestrogen metabolism in healthy post-menopausal women. *J Nutr.* 139 (5), 939-944.
- Jeannette Vera, Jorge Castro, Alberto Gonzalez, and Alejandra Moenne., 2011. Seaweed Polysaccharides and Derived Oligosaccharides Stimulate Defense Responses and Protection Against Pathogens in Plants. *Mar Drugs.* 9(12), 2514–2525.
- Jensen, A., 1972. The nutritive value of seaweed meal for domestic animals. *Proc. int. Seaweed Symp.* 7: 7–14.
- Jiao, G.; Yu, G.; Zhang, J.; Ewart, H.S., 2011. Chemical Structures and Bioactivities of Sulfated Polysaccharides from Marine Algae. *Mar. Drugs*, 9(2), 196-223.
- Johnson, N., 1990. Seaweed and its synthetic analogues in obstetrics and gynaecology 450BC-1990AD. *J R Soc Med.* 83 (6), 387-389.
- Joy L Frestedt, Melanie Walsh, Michael A Kuskowski and John L Zenk., 2008. A natural mineral supplement provides relief from knee osteoarthritis symptoms: a randomized controlled pilot trial, *Nutr J.* 7(9).
- K. Abe, T. Sakamoto, S. F. Sasaki and K. Nisizawa., 1973. In vivo studies on the

- synthesis of alginic acid in *Ishige okamurai*, *Botanica marina*, 16,(4), 229–234.
- K. Funaki and Y. Kojima. 1951. Studies on the properties of agar-agar from *Gracilaria confervoides*. Part-3, *Bull.Jap. Soc. Sci. Fish.* 16, 401-404
- K. Mikami., M. Hosokawa., 2013. Biosynthetic pathway and health benefits of fucoxanthin, an algae-specific xanthophyll in brown seaweeds, *Int. j. mol. sci.* 14 (7), 13763-13781.
- K. Sivakumar and T.Thinakaran.,2013. Antifungal activity of certain seaweeds from puthumadam coast ,*International Journal of Research and Reviews in Pharmacy and Applied science.* 3(3) 341-350.
- Kanjana K., Radtanatip T., Asuvapongpatana S., Withyachumnarnkul B., Wongprasert K., 2011. Solvent extracts of the red seaweed *gracilaria fisheri* prevent *Vibrio harveyi* infections in the black tiger shrimp *Penaeus monodon*. *Fish & shellfish immunology*, 30(1), 389–396.
- Kasi Pandima Devi., Natarajan Suganthy., Periyannaina Kesika., Shanmugaiathevar Karutha Pandian., 2008. Bioprotective properties of seaweeds: In vitro evaluation of antioxidant activity and antimicrobial activity against food borne bacteria in relation to polyphenolic content, *BMC Complementary and Alternative Medicine* 8:38. doi:10.1186/1472-6882-8-38.
- Krotkiewski M, Aurell M, Holm G, Grimby G, Szczepanik J., 1991.Effects of a sodium-potassium ion-exchanging seaweed preparation in mild hypertension, *Am J Hypertens.* 4, 483-488.
- Kusaykin M, Bakunina I, Sova V, Ermakova S, Kuznetsova T, Besednova N, Zaporozhets T, Zvyagintseva T., 2008. Structure, biological activity, and enzymatic transformation of fucoidans from the brown seaweeds, *Biotechnol. J.*, 3, (7), 904–915.
- Lalitesh Chaudhary, Priya Pradhan, Nishant Soni, Pushpendra Singh, Archana Tiwari., 2014. Algae as a Feedstock for Bioethanol Production: New Entrance in Biofuel World. *International Journal of ChemTech Research.* 6(2), 1381-1389
- Laura Giacinti, Pier Paolo Claudio, Massimo Lopez and Antonio Giordano., 2006.Epigenetic information and estrogen receptor alpha expression in breast cancer, *The Oncologist* 11 (1) 1–8.
- Lee JB, Hayashi K, Hashimoto M, Nakano T, Hayashi T., 2004. Antiviral fucoidan from sporophyll of *Undariapinnatifida* (Mekabu). *Chem. Pharm. Bull.*, 52, (9), 1091–1094.
- Leilei Ge, Peng Wang and Haijin Mou., 2011. Study on saccharification techniques of seaweed wastes for the transformation of ethanol. *Renewable Energy*, 36(1), 84-89
- Liu JN, Yoshida Y, Wang MQ, Okai Y, Yamashita U., 1997. B cell stimulating activity of seaweed extracts, *Int J Immunopharmacol.* 19 (3), 135-142.
- Liu, L, Heinrich, M, Myers, SP and Dworjanyn., 2012, Towards a better understanding of medicinal uses of the brown seaweed genus *Sargassum* in traditional Chinese medicine: a phytochemical and pharmacological review', *Journal of Ethnopharmacology*, 142, 591-619.
- M Casas-Valdez and E.A. Maina., 2006. *Sargassum* (Sargassaceae): una alternativa tropical para la alimentacion de grandos caprino, *Rev Biol Trop.* 54, 83–92.
- M. J. Abad., L. M. Bedoya., P. Bermejo.,2011. Marine Compounds and their Antimicrobial Activities, *FORMATEX 2011*, Science against microbial pathogens: communicating current research and technological advances.
- M. Povolny, 1981. The effect of the steeping of peat-cellulose flowerpots (Jiffypots) in extracts of seaweeds on the quality of tomato seedlings, *Proc Int Seaweed Symp.* 8, 730–733.
- M. Tutor-Ale, J.D. Mikkelsen, A.S. Meyer., 2011. Important Determinants for Fucoidan Bioactivity: A Critical Review of Structure-Function Relations and Extraction Methods for Fucose-Containing Sulfated Polysaccharides from Brown Seaweeds, *Marine Drugs*, 9, (10) 2106-2130.
- Mahnaz Mahdavi, Farideh Namvar, Mansor Bin Ahmad and Rosfarizan Mohamad., 2013. Green Biosynthesis and Characterization of Magnetic Iron Oxide (Fe₃O₄) Nanoparticles using Seaweed (*Sargassum muticum*) aqueous Extract, *Molecules.* 18 (5), 5954-5964.
- Marcel Tutor Ale, Jørn D. Mikkelsen, and Anne S. Meyer., 2011. Important determinants for

- fucoidan bioactivity: a critical review of structure-function relations and extraction methods for fucose-containing sulfated polysaccharides from brown seaweeds, *Marine drugs* 9 (10) 2106-2130.
- Mi-Jeong Ahn, Kee-Dong Yoon, Chul Young Kim, So-Young Min, Yong-ung Kim, Hyun Jin Kim, Jeong Ha Kim, Cha-Gyun Shin, Chong-Kyo Lee, Tae Gyun Kim, Seung Hee Kim, Hoon Huh and Jinwoong Kim., 2002. Inhibition of HIV-1 reverse transcriptase and HIV-1 integrase and antiviral activity of Korean seaweed extracts. 14, 325-329
- Moe, S. T., Draget, K. I., Skjak-Braek, G.; Smidsrod, O. 1995. Alginates. In *Food Polysaccharides and Their Applications*, ed. A.M. Stephen, Marcel Dekker, New York. 245-286.
- Mohamed A.A. Radwan, Eman A. Ragab, Nermien M. Sabry, Siham M. El-Shenawy., 2007. Synthesis and biological evaluation of new 3-substituted indole derivatives as potential anti-inflammatory and analgesic agents. *Bioorganic & Medicinal Chemistry*. 15(11), 3832-3841
- Murty U.S., Banerjee A.K. 2012. Seaweeds: The wealth of oceans. In: Kim S.-K., editor. *Handbook of Marine Macroalgae*. John Wiley & Sons, Ltd.; West Sussex, UK. 36-44.
- Murugaiyan, K., Narasimman, S., 2012. Elemental composition of *Sargassum longifolium* and *Turbinaria conoides* from Pamban Coast. *International Journal of Research in Biological Sciences*. 2(4), 137-140.
- Muruleedhara N. Byappanahalli, Dawn A. Shively, Meredith B. Nevers, Michael J. Sadowsky, Richard L. Whitman., 2003. Growth and survival of *Escherichia coli* and *Enterococci* populations in the macro-alga *Cladophora* (chlorophyta), *Fems microbiol. Ecol.* 46, 203-211.
- N. Neveux, A.K.L. Yuen, C. Jazrawi, M. Magnusson, B.S. Haynes, A.F. Masters, A. Montoya, N.A. Paul, T. Maschmeyer, R. de Nys., 2014. Biocrude yield and productivity from the hydrothermal liquefaction of marine and freshwater green macroalgae. *Bioresource Technology* 155C, 334-341.
- Nednaldo Dantas-Santos., Dayanne Lopes Gomes., Leandro Silva Costa., Sara Lima Cordeiro., Mariana Santos Santana Pereira Costa., Edvaldo Silva Trindade., Célia Regina Chavichiolo Franco., Kátia Castanho Scortecci., Edda Lisboa Leite., Hugo Alexandre Oliveira Rocha., 2012. Freshwater plants synthesize sulfated polysaccharides: Heterogalactans from water hyacinth (*Eicchorniacrassipes*), *Int. j. mol. sci.* 13 (1) 961-976.
- Nicolantonio D'Orazio . Eugenio Gemello., Maria Alessandra Gammone., Massimo de Girolamo., Cristiana Ficoneri., Graziano Riccioni., 2012. Fucoxantin: a treasure from the sea, *Mar. Drugs* .10(3), 604-616.
- Okazaki, M., Pentecost, A., Tanaka, Y. & Miyata, M., 1986. A study of calcium carbonate deposition in the genus *Padina* (Phaeophyceae, Dictyotales), *Br Phycol J.* 21, 217-224.
- P. Siamopoulou, A. Bimplakis, D. Iliopoulou *et al.*, 2004. Diterpenes from the brown algae *Dictyota dichotoma* and *Dictyota linearis*. *Phytochemistry*, 65(14). 2025-2030, Percival E., 1978. Do the polysaccharides of brown and red seaweeds ignore taxonomy? In *Modern approaches to taxonomy of red and brown algae* (Irvine, D. E. G. and Price, J. H., editors) *Systematics Association Special Volume No. 10*. Academic Press, London and New York. 47-62.
- Ponce, N. M., Pujol, C. A., Damonte, E. B., Flores, M. L., and Stortz, C. A., 2003. Fucoindans from the brown seaweed *Adenocystis utricularis*: Extraction methods, antiviral activity and structural studies. *Carbohydrate Research*. 338, 153-165.
- Ponnanikajamideen, C. Malarkodi, M. Malini and S. Rajesh kumar., 2014. Explore the antimicrobial potential from organic solvents of brown seaweed extracts (*Sargassum longifolium*) alleviating to pharmaceuticals, *International Journal of Pharmacognosy*. 1(1), 82-89.
- Pramanick, B., Brahmachari, K. and Ghosh, A., 2013. Effect of Seaweed Saps on Growth and Yield Improvement of Green Gram. *African Journal of Agricultural Research*, 8, 1180-1186.
- Quatrano RS and Stevens PT., 1976. Cell wall assembly in *Fucus zygotes*. Characterisation of the polysaccharide components, *Planta* 58(2), 224-231.
- Queiroz KC, Medeiros VP, Queiroz LS, Abreu LR, Rocha HA, Ferreira CV, Jucá MB,

- Aoyama H, Leite EL., 2008. Inhibition of reverse transcriptase activity of HIV by polysaccharides of brown algae. *Biomed Pharmacother.* 62(5), 303-307.
- Rinaudo M. 2007. Seaweed polysaccharides. In *Comprehensive Glycoscience. From Chemistry To Systems Biology* (Kamerling JP, ed). 2, 691–735.
- Rioux LE, Turgeon SL, Beaulieu M., 2010. Structural characterization of laminaran and galactofucan extracted from the brown seaweed *Saccharina longicruris*, *Phytochemistry.* 71(13) 1586-95.
- Rioux, L.E, Turgeon S.L, Beaulieu M., 2007. Characterization of polysaccharides extracted from brown seaweeds. *Carbohydrate Polymers* 69 (3), 530–537.
- Rocio Maceiras, Mónica Rodríguez, Angeles Cancela, Santiago Urréjola, Angel Sánchez., 2011. Raw material for biodiesel production, *Applied energy.* 88(10), 3318-3323.
- Rodrigues Ely, Tilvi Supriya, C.G. Naik., 2004. Antimicrobial activity of marine organisms collected off the coast of south east India, *Journal of experimental marine biology and ecology.* 309, 121– 127.
- Romanos MTV, Andrada-Serpa MJ, Santos MGM, Ribeiro ACF, Yoneshigue-Valentin Y, Costa SS & Wigg MD., 2002. Inhibitory effect of extracts of Brazilian marine algae on human T-cell lymphotropic virus type 1 (HTLV-1)-induced syncytium formation in vitro. *Cancer Investigation.* 20, 46–54.
- Ryu MJ, Kim AD, Kang KA, Chung HS, Kim HS, Suh IS, Chang WY, Hyun JW., 2013. The green algae *Ulva fasciata* delile extract induces apoptotic cell death in human colon cancer cells, *In vitro cell. dev. biol. anim.* 49 (1) 74–81.
- S. Horie, S. Tsutsumi, Y. Takada, and J. Kimura., 2008. Antibacterial quinine metabolites from the brown alga, *Sargassum sagamianum*, *Bulletin of the Chemical Society of Japan.* 81(9), 1125–1130
- S. Nagarajan and K.A. Kuppusamy., 2013. Extracellular synthesis of zinc oxide nanoparticle using seaweeds of gulf of Mannar, India. *Journal of Nanobiotechnology* 11 () 39.
- S.J. Hornl M Aasen, K. Ostgaard., 2000. Ethanol production from seaweed extract, *Journal of Industrial Microbiology & Biotechnology* 25 (5) 249-254.
- Schramm.W., Editors Guiry M.D; Blunden G. 1991. Seaweeds for waste water treatment and recycling of nutrients. Book : *Seaweed resources in Europe: uses and potential.* 149-168.
- Shan BE, Yoshida Y, Kuroda E and Yamashita., 1999. Immunomodulating activity of seaweed extract on human lymphocytes in vitro, *Int J Immunopharmacol.* 21(1), 59-70.
- Shankar SS, Rai A, Ahmad A, Sastry M., 2004. Rapid synthesis of Au, Ag, and bimetallic Au core Ag shell nanoparticles using *Neem* (*Azadirachta indica*) leaf broth, *J. Colloid Interface Sci.* 275 (2), 496–502.
- Shanmugam Rajeshkumar, Chelladurai Malarkodi, Gnanadhas Gnanajobitha, Kanniah Paulkumar, Mahendran Vanaja, Chellapandian Kannan and Gurusamy Annadurai., 2013. Seaweed-mediated synthesis of gold nanoparticles using *Turbinaria conoides* and its characterization, *Journal of Nanostructure in Chemistry.* 3 (1) 44.
- Shilpi Gupta, Nissreen Abu-Ghannam., 2011. Bioactive potential and possible health effects of edible brown seaweeds, *Trends in food science & technology* 22 (6) 315–326.
- Shota Sakai, Tatsuya Sugawara, Kiminori Matsubara, and Takashi Hirata., 2009. Inhibitory Effect of Carotenoids on the Degranulation of Mast Cells via Suppression of Antigen-induced Aggregation of High Affinity IgE Receptors. *J Biol Chem.* 284(41), 28172–28179.
- Song JY and Kim BS., 2009. Rapid biological synthesis of silver nanoparticles using plant leaf extracts, *Bioprocess Biosyst. Eng* 32 (1), 79–84.
- Subba Rao, P. V., Vaibhav, A., Mantri., 2006. Indian seaweed resources and sustainable utilization: Scenario at the dawn of a new century. *Current Science* 91 (2), 164-174.
- Suhaila Mohamed, Siti Nadia Hashim, Hafeedza Abdul Rahman., 2012. Seaweeds: A sustainable functional food for complementary and alternative therapy, *Trends in food science & technology* 23 (2) 83–96.
- Sung-Myung Kang., Seung-Hong Lee., Soo-Jin Heo., Kil-Nam Kim., You-Jin Jeon., 2011

- Evaluation of antioxidant properties of a new compound, pyrogallol-phloroglucinol-6,6'-bieckol isolated from brown algae, *Ecklonia cav.* *Nutr Res Pract.* 5(6),495–502.
- Susan Løvstad Holdt., Stefan Kraan., 2010. Bioactive compounds in seaweed: functional food applications and legislation. *Jour of app phyco.* 23 (3), 543-597.
- Suto, S., 1971. Variation in species characters of *Porphyra* under culture condition. Seminar on the Contributions of Culture, Laboratory, Field and Life History Studies to the Systematics of Benthic Marine Algae of the Pacific Abstracts. Jap.-U.S. Comm. on Sci. Coop, 28-29.
- T. Matsui, Y. Koike., 2010. Methane fermentation of a mixture of seaweed and milk at a pilot-scale plant, *Journal of Biosciences and Bioengineering*, 110, (5), 558-63.
- T. Mise, M. Ueda, T. Yasumoto., 2011. Production of fucoxanthin-rich powder from *Cladosiphon okamuranus*. *Adv. J. Food Sci. Technol.* 3 (1), 73–76.
- T.Y. Lin and W.Z. Hassid., 1966. Hassid, Isolation of guanosine diphosphate uronic acids from a brown alga, *Fucus gardneri*, *Silva. J. Biol. Chem* 241, 3283–3293.
- Tamura, T., 1966. *Marine aquaculture*, 2nd ed. (Translated from the Japanese by M. I. Watanabe, p. C-1--A2-30; available U.S. Dep. Commer., Natl. Tech Inf. Serv., Springfield, VA, as PB194051T.)
- Taratra Andréa Fenoradosoa, Ghina Ali, Cedric Delattre, Celine Laroche, Emmanuel Petit, Anne Wadouachi, Philippe Michaud ., 2010. Extraction and characterization of an alginate from the brown seaweed *Sargassum turbinarioides*, *Journal of Applied Phycology*, 22, (2), 131-137.
- Thamaraiselvan Rengarajan, Peramaiyan Rajendran, Natarajan Nandakumar, Maruthaiveeran Periyasamy Balasubramanian and Ikuo Nishigaki., 2013. Cancer preventive efficacy of marine carotenoid fucoxanthin: cell cycle arrest and apoptosis, *Nutrients* 5 (12) 4978-4989.
- Thomas Potts, Jianjun Du, Michelle Paul, Peter May, Robert Beitle and Jamie Hestekin., 2012. The production of butanol from Jamaica bay macro algae, *Environmental Progress & Sustainable Energy*, 31 (1), 29–36.
- Trincherro J, Ponce NM, Córdoba OL, Flores ML, Pampuro S, Stortz CA, Salomón H, Turk G., 2009. Antiretroviral activity of fucoidans extracted from the brown seaweed *Adenocystis utricularis*, *Phytother. Res.* 23, (5), 707–712.
- V. Ivanova, M. Stancheva, D. Petrova., 2013. Fatty acid composition of black sea *Ulvarigida* and *Cystoseira crinite*, Bulgarian . *Journal of agricultural science.* 19, 42–47
- V. Rao and I.A.B. Thivy., 1960. Seaweed utilisation in India. *Proc. Symp. Algology*, ICAR, New Delhi. pp. 345-365.
- Van der Wal H, Sperber BL, Houweling-Tan B, Bakker RR, Brandenburg W, López-Contreras AM., 2013. Production of acetone, butanol, and ethanol from biomass of the green seaweed *Ulva lactuca*. *Bioresource Technology*, 128, 431-7.
- Vigneswara Rao, A., Patel KN, Shah H.N., 1965. Manufacture of agar-agar from red seaweeds. *Res. Ind.* 10, 131-133.
- Vipul D. Prajapati · Pankaj M. Maheriya · Girish K. Jani · Himanshu K. Solanki., 2014.
- Carrageenan: A natural seaweed polysaccharide and its applications. *Carbohydrate Polymers.* 105(1), 97–112.
- Vivekanand, Vincent G. H. Eijsink, Svein J. Horn., 2012. Biogas production from the brown seaweed *Saccharina latissima*: thermal pretreatment and codigestion with wheat straw, *Journal of Applied Phycology*, 24, (5), 1295-1301.
- Vo TS, Ngo DH, Ta QV, Kim SK., 2011. Marine organisms as a therapeutic source against herpes simplex virus infection, *Eur J Pharm Sci.* 44 (1-2) 11-20.
- W. Khan, U.P. Rayirath, S. Subramanian, M.N. Jithesh, P. Rayorath, M.D. Hodges, A.T. Critchley, J.S. Craigie, J. Norrie and B. Prithiviraj., 2009. Seaweed extracts as biostimulants of plant growth and development, *J Plant Growth Regul.* 27, 270–279.
- W. Stahl., H. Sies., 2012. Photoprotection by dietary carotenoids: concept, mechanism, evidence and future development, *Mol. Nutr. Food Res.* 56 (2), 287–295.
- Wang J, Zhang Q, Zhang Z, Song H, Li P., 2010. Potential antioxidant and anticoagulant capacity of low molecular weight fucoidan fractions extracted from *Laminaria japonica*, *Int. J. Biol. Macromol.* 46 (1) 6–12.

- Wang, Y., Xu, Z., Bach, S.J., McAllister, T.A., 2009. Sensitivity of *Escherichia coli* to seaweed (*Ascophyllum nodosum*) phlorotannins and terrestrial tannins, *Asian-Australian Journal Of Animal Science* 22, 238–245.
- Wei N, Quarterman J, Jin YS., 2013. Marine macroalgae: An untapped resource for producing fuels and chemicals. *Trends Biotechnol.*31, (2), 70–7.
- Westemeier, R., Rivera, P. J., Gómez, L., Cultivo de *Gracilaria chilensis* Bird, McLachlan Oliveira., 1991. En la zona intermareal y submareal del estuario Cariquilda, Maullín, Chile. *Rev. Chil. Hist. Nat* 64, 307–321.
- Yamamoto, I., M. Takahashi, E. Tamura, H. Maruyama & H. Mori, 1984a. Antitumor activity of edible marine algae: Effect of fucoidan fractions prepared from edible seaweeds against L-1210 leukemia. *Proc. int. Seaweed Symp.* 11: 145–148.
- Yamamoto, I., T. Nagumo, K. Yagi, H. Tominaga & M. Aoki, 1974. Antitumor effect of seaweeds I. Antitumor effect of extracts from *Sargassum* and *Laminaria*. *Jpn. J. exp. Med.* 44: 543–546.
- Yamamoto, I., T. Nagumo, M. Takahashi, M. Fujihara, Y. Suzuki & N. Iizima, 1981. Antitumor effect of seaweeds III. Antitumor effect of an extract from *Sargassum kjellmanianum*. *Jpn. J. exp. Med.* 51: 187–189
- Yan X., Chuda Y., Suzuki M., Nagata T., 1999. Fucoxanthin as the major antioxidant in *Hijikia fusiformis*, a common edible seaweed, *Biosci. Biotechnol. Biochem.* 63 (3) 605–607.
- Yong-Fang HU, Yong-Jie WU., 2001. Effect of recombinant human basic fibroblast growth factor on acute inflammation in mice and rats. *Acta Pharmacol Sin.* 22, 357–379.
- Yukio Yamori MD, Ayako Miura MS and Kazuhiko Taira., 2001. Implications from and for food cultures for cardiovascular diseases: Japanese food, particularly Okinawan diets, *Asia Pac J Clin Nutr.* 10 (2), 144-145.