


**Alliance**

(<https://www.aquaculturealliance.org>)

---



Health &  
Welfare

---

# Seaweed products find varied commercial, medical applications

1 August 2005

By Dinabandhu Sahoo and Charles Yarish

## Seaweeds have value as soil additives and fertilizers

People have been eating fresh and sun-dried seaweeds for health purposes in oriental countries for thousands of years. However, in recent years, many new compounds with medicinal values have been identified in different types of seaweeds. Interestingly, most of these pharmaceutical compounds and nutraceuticals come from traditionally edible seaweeds like nori, kombu, wakame and hiziki.

## Diverse applications

The annual value of cultivated, managed and wild-harvested seaweeds exceeds \$6.0 billion, with 87 percent of that amount derived from aquaculture. With worldwide production of approximately 14 billion sheets, the most valued of the maricultured seaweeds is the red alga (*Porphyra*) popularly known as "nori" in Japan. It is a source of food for humans throughout the world, although it is primarily cultivated in China, Japan, and South Korea.

Other economically important edible seaweeds include *Laminaria* (kombu), *Undaria* (wakame), *Hizikia* (hiziki) and *Caulerpa* (sea grapes). Additional seaweed species are industrial sources of carrageenans, alginates and agars. These important polysaccharides are used in the food, textile, paint, biotechnological and biomedical industries.

Seaweeds have significant value in agriculture as soil additives, fertilizers and seaweed meals.



The most valued cultured seaweed is the red alga *Porphyra*, a high-fiber, high-protein product used in a variety of foods throughout the world.

There is also renewed interest in using seaweeds as biological nutrient removal systems in the integrated aquaculture of finfish and crustaceans. The bioremediation potential of several *Porphyra*, *Gracilaria* and *Laminaria* species has been assessed for open water and land-based systems.

## Medical uses

Many seaweeds and their polysaccharides have demonstrated antibacterial, antiviral, and antitumor effects. Recently, some seaweeds have shown antioxidant and anti-HIV potential. The sulfated polysaccharide Carraguard, now in clinical trials in Africa, appears to be a safe and cost-effective microbiocide that may reduce the transmission of HIV and possibly other sexually transmitted infections, according to the Population Council.

*Porphyra* is rich in dietary fiber, contains 25 to 50 percent protein, more vitamin C than oranges, and trace minerals. Seventeen types of free amino acids have been reported in *Porphyra*. Its very high taurine levels have been associated with reducing levels of cholesterol. *Porphyra* and *Undaria* are promising seaweeds for preventing cardiovascular disease.

Consumption of *Undaria pinnatifida* can decrease blood pressure in humans. *Enteromorpha*, *Monostroma* and *Caulerpa* are other edible seaweeds with several health benefits. Recent epidemiological studies suggested a link between the consumption of seaweeds and a reduction in prostate, colorectal, and breast cancers. A variety of antioxidant compounds including polyphenols have been identified in several brown seaweeds.

## Antiviral, antitumor

The SeaHealth project was recently launched in France to demonstrate the benefits of seaweeds' antioxidant substances to prevent atherosclerosis and cancer. Antioxidant properties have been reported from *Grateloupia filicina*, an edible red alga called limu in Hawaiian culture that is widely eaten in Hawaii, Japan, Korea, and other countries.



Cultured Kappaphycus is an important source for the carrageenan industry.

Sulphated polysaccharides from red algae inhibit viral action. Both the Herpes virus (containing DNA) and HIV (containing RNA) are inhibited by sulphated polysaccharides. Aqueous extracts of many red seaweeds are active against retroviruses. Carrageenan is cointernalized into infected cells with Herpes Simplex Virus, thus inhibiting the growth of the virus.

Many marine algae have shown antitumor activities. Significant activity against *E. coli* was found in the brown algae *Scytosiphon lomentaria* (69.8 percent inhibition), *Lessonia nigrescens* (60 percent), *Laminaria japonica* (57.6 percent) and *Sargassum ringgoldainum* (46.5 percent). Five brown and four red algae have shown appreciable antitumor properties against meth-A-fibrosarcoma.

## Other uses

Polysaccharides from brown and red seaweeds have several other uses. Earlier, alginates were used mainly to increase the viscosity of liquids and for dental filling. Today, alginates are being tailor-made by chemical selection or enzymatic treatment. Alginates require calcium, potassium, or sodium for gelation and have limited ranges of textures. Propylene glycol alginate solutions suspend particulates and are used as secondary emulsifiers and flow modifiers for salad dressings and fruit beverages.

Alginates are also used to encapsulate chemical, microbial, plant, and animal cells that are subsequently used as desired metabolite producers. Consumption of sodium alginates and low-molecular sodium alginates reduces cholesterol levels. Brown algal polysaccharides have anticoagulant, antiviral, and anti-inflammatory actions. In clinical trials, simple extracts of *Undaria pinnatifida* have been potent inhibitors of HIV.

## Culture continues

A few companies have successfully demonstrated tank cultivation of *Porphyra* for the production of specific nutraceuticals. Canada-based Ocean Nutrition and Nori Tech, an Israeli company, are developing tailor-made *Porphyra* to meet specific requirements. In India, Pepsico started large-scale cultivation of carrageenophyte seaweed.

With uses for seaweed continuing to expand, the demand for various seaweeds and their products will likely increase in the near future, thereby supporting the need for wider development of pharmacological aquaculture.

*(Editor's Note: This article was originally published in the August 2005 print edition of the Global Aquaculture Advocate.)*

## Authors

---



### DINABANDHU SAHOO

Marine Biotechnology Laboratory  
Department of Botany  
University of Delhi  
Delhi, India



### CHARLES YARISH

Department of Ecology and Evolutionary Biology  
University of Connecticut  
1 University Place  
Stamford, Connecticut 06901-2315 USA

[charles.yarish@uconn.edu](mailto:charles.yarish@uconn.edu) (mailto:charles.yarish@uconn.edu)

All rights reserved.